

# TENNESSEE PERINATAL CARE SYSTEM

## GUIDELINES FOR TRANSPORTATION

(Fifth Edition)



2006

Tennessee Department of Health  
Women's Health & Genetics

Phil Bredesen  
Governor

Kenneth S. Robinson, M.D.  
Commissioner

**TENNESSEE PERINATAL CARE SYSTEM  
GUIDELINES FOR TRANSPORTATION**

**(Fifth Edition)**

**Prepared by the  
Subcommittee on Perinatal Transportation  
of the  
Perinatal Advisory Committee**

**2006**

This document may be accessed electronically at the following:

<http://www.state.tn.us/health/>

Click on Forms and Publications and then Emergency Medical Services or Women's Health

# TRANSPORTATION GUIDELINES WORKGROUP

April 2006

## **Members**

Cheryl W. Major, R.N.C., B.S.N., Chairperson, Nashville

C. David Adair, M.D., Chattanooga and Knoxville

Kitty Cashion, R.N., B.C., M.S.N., Memphis

W. Michael DeVoe, M.D., Johnson City

Susan Drummond, R.N., M.S.N., Nashville

Ann Everson, R.N., Chattanooga

Mark S. Gaylord, M.D., Knoxville

Connie Graves, M.D., Nashville

James C. Gray, M.D., Cookeville

Rita R. Hillhouse, R.N.C., Knoxville

Sami Ismail, M.D., Nashville

Patti Jacobs, R.N., Johnson City

Steve Jones, E.M.T., Lafayette

Richard Land, E.M.T.-P, Nashville

Terry A. Longhofer, R.N.C., N.N.P., Knoxville

Chris Martin, M.D., Kingsport

Mitchell Mask, M.S., R.R.T., N.P.S., Memphis

Lisa McClane, R.N.C., M.S.N., C.N.S., Chattanooga

Margaret Rush, M.D., Nashville

Patti Scott, M.S.N., R.N.C., A.P.N.- N.N.P., Nashville

Ajay Talati, M.D., Memphis

Judy Wallace, R.N.C, Johnson City

Terri Woodruff, R.N., M.S., C.P.N.P., Chattanooga

# PERINATAL ADVISORY COMMITTEE

March 2006

## Members

Ruth Hagstrom, M.D., Chairman, Nashville

David Adair, M.D., Chattanooga

Judy Ascher, M.D., Nashville

Des Bharti, M.D., Johnson City

W. Michael DeVoe, M.D., Johnson City

Ramasubbareddy Dhanireddy, M.D., Memphis

Connie Graves, M.D., Nashville

J. Chris Graves, M.D., Chattanooga

Rita R. Hillhouse, R.N.C., Knoxville

Lizbeth Kennedy, M.D., Chattanooga

Vichien Lorch, M.D., Knoxville

Cheryl W. Major, R.N.C., B.S.N., Nashville

Norman L. Meyer, M.D., Ph.D., Memphis

Martin Olsen, M.D., Johnson City

Marilyn Robinson, M.D., Memphis

Jorge Rojas, M.D., Nashville

Donald W. Tansil, M.D., Cookeville

Lenita H. Thibault, M.D., Kingsport

Dennis L. Vonderfecht, Johnson City

Janice Whitty, M.D., Nashville

## **TABLE OF CONTENTS**

## TABLE OF CONTENTS

	<u>Page</u>
<b>PREFACE</b> .....	1
<b>INTRODUCTION</b> .....	2
<b>LEVEL I FACILITIES</b>	
<b>Maternal Transport</b>	
Indications for Maternal Consultation and/or Transport .....	4
Maternal Referral Process.....	6
Maternal Transport Personnel.....	8
Maternal Transport Modality.....	9
Maternal Transport Equipment.....	10
Maternal Referral Documentation.....	12
Evaluation of Perinatal Referral Process.....	13
<b>Neonatal Transport</b>	
Indications for Neonatal Consultation and/or Transport .....	14
Neonatal Referral Process .....	15
Neonatal Transport Personnel .....	17
Neonatal Referral Documentation .....	18
Evaluation of Neonatal Referral Process .....	19
<b>LEVEL II-A FACILITIES</b>	
<b>Maternal Transport</b>	
Indications for Maternal Consultation and/or Transport .....	20
Maternal Referral Process.....	22
Maternal Transport Personnel.....	24
Maternal Transport Modality.....	25
Maternal Transport Equipment.....	26
Maternal Referral Documentation.....	28
Evaluation of Perinatal Referral Process.....	29
<b>Neonatal Transport</b>	
Indications for Neonatal Consultation and/or Transport .....	30
Neonatal Referral Process .....	31
Neonatal Transport Personnel .....	33
Neonatal Transport Modality .....	34
Neonatal Referral Documentation .....	35
Evaluation of Neonatal Referral Process.....	36
<b>LEVEL II-B FACILITIES</b>	
<b>Maternal Transport</b>	
Indications for Maternal Consultation and/or Transport .....	37
Maternal Referral Process.....	39

Maternal Transport Personnel.....	41
Maternal Transport Modality.....	42
Maternal Transport Equipment.....	43
Maternal Referral Documentation.....	45
Evaluation of Perinatal Referral Process.....	46

**Neonatal Specialty Transport Modality**

Indications for Neonatal Consultation and/or Transport.....	47
Neonatal Referral Process.....	48
Neonatal Transport Personnel.....	50
Neonatal Specialty Transport Modality.....	51
Neonatal Transport Equipment.....	53
Neonatal Referral Documentation.....	57
Evaluation of Neonatal Referral Process.....	58

**LEVEL III FACILITIES**

**Maternal Transport**

Indications for Maternal Consultation and/or Transport.....	59
Maternal Referral Process.....	60
Maternal Transport Personnel.....	62
Maternal Transport Modality.....	63
Maternal Transport Equipment.....	64
Maternal Referral Documentation.....	66
Evaluation of Perinatal Referral Process.....	67

**Neonatal Specialty Transport Modality**

Indications for Neonatal Consultation and/or Transport.....	68
Neonatal Referral Process.....	69
Neonatal Transport Personnel.....	71
Neonatal Specialty Transport Modality.....	72
Neonatal Transport Equipment.....	74
Neonatal Referral Documentation.....	78
Evaluation of Neonatal Referral Process.....	79

<b>RETURN TRANSPORT.....</b>	<b>80</b>
------------------------------	-----------

**APPENDICES**

- I. Educational Objectives for Neonatal Transport Nurses
- II. Education Requirements for Emergency Medical Services Personnel
- III. Oxygen Cylinder Usage During Transport
- IV. An Example of a Neonatal Return Transport Worksheet
- V. An Example of a Neonatal Return Transport Follow-up Form
- VI. COBRA/EMTALA Statute: 42 USC 1395
- VII. Neonatal Resuscitation Equipment and Supplies
- VIII. BLS / ALS Equipment and Supplies

# **PREFACE**

# **TENNESSEE PERINATAL CARE SYSTEM**

## **GUIDELINES FOR TRANSPORTATION**

### **PREFACE**

This manual is a revision of the Guidelines for Transportation in the State of Tennessee that was originally published in September 1979. These guidelines are written in response to the recommendation of the Perinatal Advisory Committee and are developed to accomplish improvement in the overall quality of maternal-newborn transportation in the state. Prepared by the Subcommittee on Perinatal Transportation, and adopted by its parent committee, the Perinatal Advisory Committee, this manual has been carefully considered by representatives from a broad spectrum of the health care delivery disciplines from throughout the state. This manual provides specific guidelines regarding procedures, staffing patterns, and equipment for the transport of high-risk mothers and neonates. It is hoped that physicians, nurses, respiratory therapists, and other health care providers involved with such transportation will make reasonable efforts to attain the guidelines described herein.

In order to insure the contemporary pertinence of these guidelines, the Perinatal Advisory Committee has limited its approval to a period that is no longer than five years from the date of approval by the Commissioner of the Department of Health. A complete review will thus be mandatory at that time or sooner if deemed appropriate.

# **INTRODUCTION**

## INTRODUCTION

The transport of pregnant women and newborn infants between hospitals is recognized as an essential component of regionalized perinatal care. National experience suggests that perinatal outcome for high-risk infants transported before delivery (maternal transport) is improved over that for high-risk infants transported after birth (neonatal transport). Tennessee experience also supports this. The Subcommittee on Perinatal Transportation of the Perinatal Advisory Committee, therefore, strongly urges that maternal transport to an appropriate referral center be considered, especially in those pregnancies in which there is a high probability of neonatal transport following delivery (Resolution passed by the Perinatal Advisory Committee on March 31, 1982).

Safe transport of the perinatal patient requires skilled personnel, appropriate equipment, and effective communication between the hospital facilities of the region. The purpose of this manual is to present specific guidelines to physicians, nurses, and other health care providers involved with maternal-neonatal transport, so that such a transfer may be conducted in the most optimal manner.

This manual is divided into four main sections: Level I Facilities, Level II-A Facilities, Level II-B Facilities, and Level III Facilities. The guidelines within each section are organized in the subsections highlighted below. In each of these subsections, maternal transport discussion precedes neonatal discussion.

1. Indications for Consultation and/or Transport

The circumstances in which patients are transported vary according to the level of care of the facilities in each region. The indications highlighted in this subsection are designed to assist the health care providers in seeking consultation and/or transfer. These indications remain guidelines and will vary with individual patient needs and institutional capabilities.

2. Referral Process

The transfer of care of a perinatal patient from one care facility to another requires effective communication between facilities and a clear understanding of the responsibilities of the parties involved. The guidelines highlighted in this subsection are designed to clarify the roles of both referring and receiving facilities.

3. Transport Personnel

The level of expertise of the transport personnel needed to provide optimal transfer of pregnant women and newborn infants is variable and largely dependent on the complexity of care demanded by the individual patient. However, in order to eliminate any possibility of encountering a situation in which the demands of the transported patient exceed the care level that can be adequately provided, it is desirable to define certain minimal requirements. The guidelines highlighted in this subsection are designed to assist the health care providers in selecting appropriate transport personnel.

4. Transport Modality

The choice of transport modality is largely determined by the distance between hospitals, weather, condition of the patient, and equipment available in the vehicle to support the patient during transfer. The guidelines highlighted in this subsection are designed to assist the health care providers in selecting the appropriate transport modality.

5. Transport Equipment

A safe perinatal transport entails availability of adequate equipment for monitoring, resuscitation, and support of both mother and neonate. In addition to the equipment in the transport vehicle, essential supplies should be portable and continuously available during the transport. The equipment needs highlighted in this subsection are designed to assist the health care providers in selecting appropriate transport equipment.

6. Referral Documentation

An essential component of communication between care facilities is clear documentation and transfer of medical information. The guidelines highlighted in this subsection are designed to clarify the responsibilities of both referring and receiving facilities with respect to medical documentation.

7. Evaluation of Referral Process

The success of a regionalized perinatal care system depends on an ongoing evaluation of various aspects of the program including transport of the perinatal patient. This subsection highlights the need for evaluation of the referral process.

Optimal utilization of a regionalized perinatal care system entails early planning and return transport of patients from the referral centers to the original or local hospitals for further care. This manual ends with a section on return transport in which various aspects of this activity are summarized.

## **LEVEL I FACILITIES**

# **MATERNAL TRANSPORT**

## **Level I**

Level I units provide basic care for uncomplicated maternity and neonatal patients. All high-risk mothers and neonates must be promptly identified for referral and/or consultation for more specialized care. The Level I unit shall provide equipment and staff to care for maternity patients whose onset of labor occurs on or after the first day of the 35<sup>th</sup> week, for neonates whose birthweight is over 2000 grams, or for sick patients pending transfer to another hospital.

Planned deliveries at gestational ages below 35 weeks should be referred to a Level II-B or III facility. Although it is not always possible to prenatally anticipate the need for pediatric subspecialty services, when antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for delivery in a facility with the needed services is encouraged.

When transfer is deemed safe for the mother and fetus, transfer of mothers to a Level II-B or III unit with the specialized expertise required by the fetus after birth is encouraged. The elective delivery of an infant in a hospital without the required pediatric sub-specialty services resulting in a planned neonatal transport is discouraged.

## INDICATIONS FOR MATERNAL CONSULTATION AND/OR TRANSPORT

### I. ANTEPARTUM

#### A. Maternal History

1. previous preterm labor (<37 weeks) or low-birthweight neonate (<2500 gm)
2. previous neonate >4000 gm at term or any large-for-gestational age neonate
3. previous stillbirth, neonatal loss, or two or more abortions
4. suspected incompetence of the cervix
5. diagnosed abnormality of the genital tract
6. medical indication for termination of previous pregnancy
7. neonate who required more than routine observation or care
8. neonate with known or suspected genetic disorder
9. severe emotional problems associated with previous pregnancy or delivery
10. previous vertical or classical uterine incision
11. age <16 or advanced maternal age ( $\geq 35$  years of age at delivery)
12. prepregnancy weight <45 kg or >90 kg
13. height <150 cm

#### B. Medical/Surgical Complications

1. diabetes mellitus/endocrine disorder
2. autoimmune disorder
3. cardiac disease
4. hypertension
5. pulmonary disease
6. renal disease
7. hematologic disorder
8. neurologic disorder
9. musculoskeletal disorder
10. infection
11. nutritional disorder
12. substance use
13. malignancy
14. psychiatric disorder
15. trauma
16. surgical emergency
17. morbid obesity

#### C. Obstetric Complications

1. glucose intolerance
2. urinary tract infection
3. sexually transmitted disease
4. positive fetal fibronectin test
5. suspected ectopic pregnancy

6. suspected missed abortion
7. hyperemesis
8. exposure to teratogen
9. isoimmunization
10. persistent anemia
11. vaginal bleeding
12. preeclampsia/eclampsia
13. suspected polyhydramnios or oligohydramnios
14. preterm cervical dilatation without uterine activity
15. preterm rupture of membranes with or without uterine activity
16. rupture of membranes at term for more than 12 hours without labor and/or evidence of amnionitis or sepsis at any time
17. suspected feto-pelvic disproportion
18. inappropriate fetal growth for gestational age
19. multiple gestation
20. postterm gestation (>42 weeks)
21. fetal demise
22. known or suspected fetal anomaly
23. abnormal triple screen
24. morbid obesity

## **II. INTRAPARTUM**

- A. preterm (<37 weeks) cervical dilatation with uterine contractions
- B. abnormal presentation
- C. suspected feto-pelvic disproportion
- D. dysfunctional labor
- E. rupture of membranes at term for more than 12 hours and/or evidence of amnionitis or sepsis at any time
- F. abnormal bleeding
- G. suspected nonreassuring fetal status
- H. preeclampsia/eclampsia
- I. uterine hyperstimulation syndrome
- J. meconium in amniotic fluid
- K. multiple gestation
- L. morbid obesity

## **III. POSTPARTUM**

- A. preeclampsia/eclampsia
- B. sepsis
- C. abnormal bleeding
- D. thromboembolic disease
- E. cardiopulmonary dysfunction
- F. morbid obesity
- G. neonatal transport

## **MATERNAL REFERRAL PROCESS**

I. Maternal transport is initiated by the health care provider responsible for the patient's medical care. Maternal referral may lead to admission of the patient to the receiving hospital (inpatient transport) or to outpatient evaluation and management (outpatient transport). The guidelines for the referral process are outlined below.

### **II. INPATIENT TRANSPORT**

#### **A. Referring Center Responsibilities:**

1. The decision by the referring care provider (physician, certified nurse midwife, nurse practitioner) to request consultation is the first step in the referral process.
2. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the care provider in developing a treatment plan for stabilizing the patient before and during transport.
3. Personnel of the referring and receiving facilities must follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI.). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.
4. If the transport is done by the referring hospital, the referring physician and hospital retain responsibility until the transport team arrives with the patient at the receiving hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
5. A ground ambulance is the most appropriate vehicle for the majority of maternal transports. If an alternate form of transportation is being considered, the referring physician should discuss this alternative with the receiving physician at the time of consultation.
6. The medical record should be organized as the patient is prepared for transport and sent with the patient.
7. The composition of the transport team should be a joint decision between the referring and receiving care providers based on the condition of the mother and/or fetus.
8. To avoid unnecessary delays in the emergency room or admitting office, all referrals should be directly admitted to the receiving obstetric unit.

#### **B. Receiving Center Responsibilities:**

1. The receiving physician is responsible for the decision to accept the referring care provider's request for transport and make preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring physician in locating appropriate care.
2. Personnel of the referring and receiving facilities must follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social

Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.

3. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
4. Every patient accepted by the receiving center should be seen by a physician within 30 minutes of arrival.
5. Communication with the referring care provider should occur following admission.
6. If the patient is discharged undelivered, communication should occur prior to the time of discharge.
7. A summary of care of both mother and/or infant should be sent to the referring care provider.
8. Every effort should be made to return the patient to the care of the referring care provider as soon as possible.

### **III. OUTPATIENT REFERRAL**

#### **A. Referring Center Responsibilities:**

1. Outpatient referral should begin with a phone call from the referring care provider to the receiving physician.
2. A convenient time for evaluation of the patient at the receiving center should be arranged.

#### **B. Receiving Center Responsibilities:**

1. The referring care provider should be contacted by either telephone or letter.
2. Whenever possible, patients should continue under the care of the referring care provider.

## **MATERNAL TRANSPORT PERSONNEL**

- I.** The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and/or fetus.
- II.** The transport team members should be selected from appropriately trained, licensed health care providers.
- III.** Transport team members should have the collective expertise sufficient to provide the following, if necessary:
  - A.** monitoring of blood pressure, uterine contractions, deep tendon reflexes, and fetal heart rate
  - B.** monitoring the administration of intravenous infusions and usage of tocolytic, antihypertensive, and anticonvulsant medications
  - C.** care for a wide variety of emergency conditions including delivery and neonatal resuscitation.
- IV.** Transport team members should be oriented to the transport vehicle and usage of transport equipment. All transport team members should follow state and national standards.
- V.** In instances such as advanced labor, unstable maternal condition, or severe illness, it may become necessary for the referring care provider to accompany the patient during transport, if transport is still recommended by the receiving physician. (See Appendix VI re EMTALA)

## MATERNAL TRANSPORT MODALITY

- I. Selection of the transport modality should be a joint decision by the referring care provider and receiving physician(s) based on the condition of the mother and fetus.
- II. Maternal transport can be accomplished by private vehicle, ambulance, rotary wing aircraft (RWA), or fixed wing aircraft (FWA). The ambulance (land and air) must be licensed by the Emergency Medical Services Division of the Tennessee Department of Health or sanctioned by Tennessee statute.
- III. If care en route is necessary, an Advanced Life Support (ALS) ambulance is required for maternal transport. The description of an ALS ambulance is defined by the Tennessee Department of Health and may be located in the latest edition of the Tennessee Emergency Medical Services Statutes and Rules, located at [www.tennessee.gov](http://www.tennessee.gov) under the "Laws & Justice" heading.

## MATERNAL TRANSPORT EQUIPMENT

- I. The referring health care provider should be aware of the availability of Advanced Life Support (ALS) ambulances in the area. The required equipment and supplies for maternal and neonatal care in an ALS ambulance are defined by the Tennessee Department of Health (See Appendix VIII).
- II. Organization and maintenance of additional transport equipment is the responsibility of the transport team.
- III. Additional equipment and supplies that may be necessary should be provided by the transporting team. These include:
  - A. doppler
  - B. reflex hammer
  - C. infusion pump
  - D. oxygen masks (premature and newborn size)
  - E. neonatal resuscitation supplies and equipment (See Appendix VII)
  - F. suction catheters (#6, #8, and #10 Fr)
  - G. latex free equipment and supplies must be available
- IV. Additional medications (or therapeutic equivalents) that may be necessary should be provided by the transport team. Such medications, including but not limited to those listed below, may be given when ordered by the referring physician.
  - A. Antenatal corticosteroids to accelerate fetal lung maturity
    1. Betamethasone for injection (12 mg IM is usual dose)
    2. Dexamethasone for injection (6–10 mg IM is usual dose)
  - B. Eclampsia
    1. Magnesium Sulfate 6 gram bolus IV, then 2 grams/hr IV. Additional dose of 2 grams over 5-10 minutes for persistent seizures (repeat x 1)
    2. Calcium gluconate 1 gram IV to reverse magnesium overdose
    3. Sodium amobarbital 250 mg IV over 3 minutes (can substitute another short-acting benzodiazepine)
  - C. Oxytocics
    1. Oxytocin (Pitocin) 10 units per ampule / vial
    2. Misoprostol (Cytotec) 100 mcg tablets and pill cutter
    3. Methylergonovine (Methergine) 200 mcg ampules
    4. Carboprost (Hemabate) 250 microgram ampule / vial
  - D. Tocolytics
    1. Terbutaline Sulfate for injection (0.25 mg subcutaneously at 20 to 60 minute intervals is usual dose)
    2. Magnesium Sulfate (dosage same as for eclampsia)
  - E. Antibiotics for Group B strep prophylaxis (ref: CDC MMWR, August 16, 2002, / Vol.

51 / No. RR-11)

1. Penicillin G 5,000,000 units IV
2. Ampicillin 2 grams IV
3. Cefazolin 2 grams IV
4. Clindamycin 900 mg IV
5. Erythromycin 500 mg IV
6. Vancomycin 1 gram IV

F. Antihypertensives

1. Labetalol (Trandate) IV. Dosage is repeated and/or adjusted at 20 minute intervals according to patient response. May sequentially give 20 mg, then 40 mg, then 80 mg, then an additional 80 mg, if insufficient response to the lower doses. Maximum dosage is 220 mg–300 mg.
2. Hydralazine (Apresoline) 5-10 mg IV every 20 minutes. Maximum dosage is 30 mg.
3. Nifedipine (Procardia) 10 mg p.o. every 20 minutes. Maximum dosage is 100 mg.

## **MATERNAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both the referring and receiving centers have responsibilities to provide adequate documentation of clinical data.

### **II. REFERRING CENTER RESPONSIBILITIES**

- A. The following documents should accompany the transported patient:
  - 1. copy of complete prenatal record
  - 2. copy of current medical record (include the EMTALA form)
  - 3. copy of record of care during transport
- B. The referring center should maintain a record regarding disposition of transported mothers.

### **III. RECEIVING CENTER RESPONSIBILITIES**

- A. maintain a record of consultation/referral calls
- B. maintain a record regarding the disposition of the transported patient
- C. send a summary of care of both mother and/or infant to the referring care provider

## EVALUATION OF PERINATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk perinatal patient requires the cooperation and coordination of many skilled health care personnel. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available through the perinatal network
  - B. assisting perinatal care providers in developing their abilities to identify high-risk perinatal patients, anticipate complications, and stabilize those patients before transport
  - C. continuing quality improvement through ongoing education of perinatal care and EMS providers (See Appendices I and II).
- III. Planning of the perinatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their maternal referrals with or without the assistance of the receiving center.

# **NEONATAL TRANSPORT**

## **Level I**

Level I units provide basic care for uncomplicated maternity and neonatal patients. All high-risk mothers and neonates must be promptly identified for referral and/or consultation for more specialized care. The Level I unit shall provide equipment and staff to care for maternity patients whose onset of labor occurs on or after the first day of the 35<sup>th</sup> week, for neonates whose birthweight is over 2000 grams, or for sick patients pending transfer to another hospital.

Planned deliveries at gestational ages below 35 weeks should be referred to a Level II-B or III facility. Although it is not always possible to prenatally anticipate the need for pediatric subspecialty services, when antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for delivery in a facility with the needed services is encouraged.

When transfer is deemed safe for the mother and fetus, transfer of mothers to a Level II-B or III unit with the specialized expertise required by the fetus after birth is encouraged. The elective delivery of an infant in a hospital without the required pediatric subspecialty services resulting in a planned neonatal transport is discouraged.

## **INDICATIONS FOR NEONATAL CONSULTATION AND/OR TRANSPORT**

- I.** Requirement for more than routine care as prescribed for normal neonates as published in the most recent edition of Guidelines for Perinatal Care (American Academy of Pediatrics and American College of Obstetricians and Gynecologists).
- II.** Gestational age <35 weeks or birthweight <2000 gm
- III.** Apgar score <3 at 1 minute, <5 at 5 minutes, and/or <7 at 10 minutes
- IV.** Need for oxygen therapy after initial resuscitation
- V.** Abnormal respirations with or without need for supplemental oxygen
- VI.** Requirement for continuous intravenous therapy >24 hours
- VII.** Suspected sepsis
- VIII.** Suspected congenital heart disease
- IX.** Neurologic disorder
- X.** Gastrointestinal disorder
- XI.** Genitourinary disorder
- XII.** Hematologic disorder
- XIII.** Musculoskeletal disorder
- XIV.** Endocrine or metabolic disorder
- XV.** Congenital malformation or suspected genetic disorder requiring further evaluation

## NEONATAL REFERRAL PROCESS

- I. Neonatal transport is initiated by the health care provider responsible for the patient's medical care. Guidelines for the referral process are outlined below.

### II. REFERRING CENTER RESPONSIBILITIES

- A. The referring health care provider's decision to request consultation is the first step in the referral process.
- B. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the referring health care provider in developing a treatment plan for stabilizing the patient before transport.
- C. A discussion between the referring health care provider and receiving physician regarding the neonate will result in one of three possible dispositions:
  1. Required neonatal care can be provided at the referring center. The receiving physician under these circumstances has only a consultative role.
  2. The neonate requires further observation, investigation, or other preparation before possible transport. Continued contact between the providers is necessary.
  3. Transport of the neonate is necessary. The optimal time, mode of transfer, transport personnel, and additional information regarding the neonate should be discussed. The process of stabilization of the neonate at the referring center should be reviewed and documented.
- D. If the neonate is transported by the referring center, the referring health care provider is responsible for the patient until arrival at the receiving center.
- E. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- F. The medical record should be organized as the patient is prepared for transport and sent with the patient.
- G. Consent forms to authorize transfer must be obtained.
- H. Parents should be encouraged to see and touch the infant. Photographs of the infant should be provided to parents, if culturally appropriate.
- I. Appropriate maternal/neonatal identification should be in place before transport.
- J. All referrals should be directly admitted to the receiving unit to avoid unnecessary delays in the emergency room.

### III. RECEIVING CENTER RESPONSIBILITIES

- A. The receiving physician is responsible for the decision to accept the referring health care provider's request for transport and make preparations at the receiving center.

- If unable to accept the transport, assistance will be provided to the referring health care provider in locating appropriate care.
- B. If the neonate is transported by the referring center, full responsibility begins with admission to the receiving center.
  - C. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
  - D. If a delay in transport occurs, a repeat telephone call should be made for further assessment and advice.
  - E. On arrival of the transport team, assessment and further stabilization of the neonate should be done in collaboration with the referring staff.
  - F. The transport team must verify proper identification of the neonate before transport.
  - G. Following stabilization of the neonate, the transport team should communicate with the parents to assure their understanding of the infant's condition, possible course, and therapeutic intervention that has been undertaken or anticipated. Written information about the receiving center should be provided to the parents.
  - H. Consent forms to authorize transfer, treatment, and admission to the receiving hospital must be obtained.
  - I. Parents should be encouraged to see and touch the infant before departure. Photographs of the infant should be provided to parents, if culturally appropriate.
  - J. Before departure, the transport team should communicate with receiving center physician regarding the neonate's history, current status, and planned management during transport.
  - K. Before arrival at the receiving center, the transport team should communicate the status of the neonate and anticipated needs on admission.
  - L. A telephone call should be made to the parents shortly following admission of the neonate.
  - M. Within 24 hours of admission, communication with referring center personnel regarding events during transport and since admission should occur.
  - N. Periodic communication with the referring health care provider should be maintained.
  - O. Consideration should be given to returning the care of the patient to the referring health care provider or primary care physician when practical and medically appropriate.
  - P. Upon discharge of the infant, a discharge summary should be sent to the referring perinatal health care providers.

## NEONATAL TRANSPORT PERSONNEL

- I. The composition of the transport team should be decided jointly by the referring health care provider and receiving physician based on the condition of the neonate. Transport by a Level I unit should be a rare event, done only after consultation with a Level III unit. In most instances neonates requiring Level II or III care should be transported by a team meeting Level III guidelines.
- II. Transport team members should be selected from appropriately trained, licensed health care providers.
- III. Transport team members should have the collective expertise sufficient to provide the following, if necessary:
  - A. observation of the newborn throughout the transport
  - B. monitoring of body temperature, respiratory status, and cardiovascular status
  - C. delivery and monitoring of oxygen therapy
  - D. initiation and maintenance of IV access and therapy
  - E. supportive care for a wide variety of emergency conditions, including positive pressure ventilation.
- IV. At least one member of the transport team should be a current Neonatal Resuscitation Program (American Academy of Pediatrics and American Heart Association) provider.
- V. Transport team members should be oriented to the transport vehicle and use of transport equipment. All transport team members should follow state and national standards.
- VI. The knowledge and skills required for a nurse to perform neonatal transport are listed in the most current edition of the Educational Objectives for Nurses, Levels I, II, III, Neonatal Transport Nurses, Tennessee Perinatal Care System, Tennessee Department of Health (Appendix I). Other suggested supplemental education includes the AAP/AHA Neonatal Resuscitation Program and The S.T.A.B.L.E. Program<sup>®</sup>, endorsed by the AAP and the March of Dimes.

## **NEONATAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both referring and receiving center personnel have responsibilities to provide adequate documentation of clinical data.
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported neonate:
    - 1. copy of complete maternal prenatal record
    - 2. copy of current maternal medical record
    - 3. copy of current neonatal medical record
    - 4. copies of any images in a format that can be reviewed by the transport team at the bedside
    - 5. copy of record of care during transport
  - B. The referring center should maintain a record regarding disposition of transferred neonates.
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported neonate
  - C. send a summary of care to the referring perinatal health care providers

## EVALUATION OF NEONATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk neonate requires the cooperation and coordination of many skilled health care personnel. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available to them through the perinatal network.
  - B. assisting neonatal care providers in developing their abilities to identify high-risk neonatal patients, anticipate complications, and stabilize those patients before transport.
  - C. continuing quality improvement through ongoing education of neonatal care and EMS providers (See Appendices I and II).
- III. Planning of the neonatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their neonatal referrals, with or without the assistance of the receiving center.

## **LEVEL II-A FACILITIES**

# **MATERNAL TRANSPORT**

## **Level II-A**

Level II-A units provide care for maternal and neonatal patients born at 34 weeks gestation or above whose courses are uncomplicated and for patients with mild obstetric and neonatal illnesses who do not require specialized services.

When antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for delivery in a facility with the needed services is encouraged.

When transport is deemed safe for the mother and fetus, transfer of mothers to a Level III unit with the specialized services required by the fetus after birth is encouraged. The elective delivery of an infant in a hospital without the required pediatric subspecialty services resulting in a planned neonatal transport is discouraged.

The nursery must have equipment and personnel to provide controlled thermal environments, hood oxygen for protracted management and assisted neonatal ventilation pending transfer to another institution for more specialized care. Obstetric and pediatric co-directors are board-certified in their respective specialties.

## INDICATIONS FOR MATERNAL CONSULTATION AND/OR TRANSPORT

### I. ANTEPARTUM

#### A. Maternal History

1. previous preterm labor (<37 weeks) or low-birthweight neonate (<2500 gm)
2. previous neonate >4000 gm at term or any large-for-gestational age neonate
3. previous stillbirth, neonatal loss, or two or more abortions
4. suspected incompetence of the cervix
5. diagnosed abnormality of the genital tract
6. medical indication for termination of previous pregnancy
7. neonate who required more than routine observation or care
8. neonate with known or suspected genetic disorder
9. severe emotional problems associated with previous pregnancy or delivery
10. previous vertical or classical uterine incision
11. age <16 or advanced maternal age ( $\geq 35$  years of age at delivery)
12. prepregnancy weight <45 kg or >90 kg
13. height <150 cm

#### B. Medical/Surgical Complications

1. diabetes mellitus/endocrine disorder
2. autoimmune disorder
3. cardiac disease
4. hypertension
5. pulmonary disease
6. renal disease
7. hematologic disorder
8. neurologic disorder
9. musculoskeletal disorder
10. infection
11. nutritional disorder
12. substance use
13. malignancy
14. psychiatric disorder
15. trauma
16. surgical emergency
17. morbid obesity

#### C. Obstetric Complications

1. glucose intolerance
2. urinary tract infection
3. sexually transmitted disease
4. positive fetal fibronectin test
5. suspected ectopic pregnancy

6. suspected missed abortion
7. hyperemesis
8. exposure to teratogen
9. isoimmunization
10. persistent anemia
11. vaginal bleeding
12. preeclampsia/eclampsia
13. suspected polyhydramnios or oligohydramnios
14. preterm cervical dilatation without uterine activity
15. preterm rupture of membranes with or without uterine activity
16. rupture of membranes at term for more than 12 hours without labor and/or evidence of amnionitis or sepsis at any time
17. suspected feto-pelvic disproportion
18. inappropriate fetal growth for gestational age
19. multiple gestation
20. postterm gestation (>42 weeks)
21. fetal demise
22. known or suspected fetal anomaly
23. abnormal triple screen
24. morbid obesity

## **II. INTRAPARTUM**

- A. preterm (<37 weeks) cervical dilatation with uterine contractions
- B. abnormal presentation
- C. suspected feto-pelvic disproportion
- D. dysfunctional labor
- E. rupture of membranes at term for more than 12 hours and/or evidence of amnionitis or sepsis at any time
- F. abnormal bleeding
- G. suspected nonreassuring fetal status
- H. preeclampsia/eclampsia
- I. uterine hyperstimulation syndrome
- J. meconium in amniotic fluid
- K. multiple gestation
- L. morbid obesity

## **III. POSTPARTUM**

- A. preeclampsia/eclampsia
- B. sepsis
- C. abnormal bleeding
- D. thromboembolic disease
- E. cardiopulmonary dysfunction
- F. morbid obesity
- G. neonatal transport

## **MATERNAL REFERRAL PROCESS**

I. Maternal transport is initiated by the health care provider responsible for the patient's medical care. Maternal referral may lead to admission of the patient to the receiving hospital (inpatient transport) or to outpatient evaluation and management (outpatient transport). The guidelines for the referral process are outlined below.

### **II. INPATIENT TRANSPORT**

#### **A. Referring Center Responsibilities:**

1. The decision by the referring care provider (physician, certified nurse midwife, nurse practitioner) to request consultation is the first step in the referral process.
2. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the care provider in developing a treatment plan for stabilizing the patient before and during transport.
3. Personnel of the referring and receiving facilities must follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.
4. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
5. A ground ambulance is the most appropriate vehicle for the majority of maternal transports. If an alternate form of transportation is being considered, the referring physician should discuss this alternative with the receiving physician at the time of consultation.
6. The medical record should be organized as the patient is prepared for transport and sent with the patient.
7. The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and fetus.
8. To avoid unnecessary delays in the emergency room or admitting office, all referrals should be directly admitted to the receiving obstetric unit.

#### **B. Receiving Center Responsibilities:**

1. The receiving physician is responsible for the decision to accept the referring care provider's request for transport and making preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring physician in locating appropriate alternative care.
2. Personnel of the referring and receiving facilities must follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social

Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.

3. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
4. Every patient accepted by the receiving center should be seen by a physician within 30 minutes of arrival.
5. Communication with the referring care provider should occur following admission.
6. If the patient is discharged undelivered, communication should occur prior to the time of discharge.
7. A summary of care of both mother and infant should be sent to the referring care provider.
8. Every effort should be made to return the patient to the care of the referring care provider as soon as possible.

### **III. OUTPATIENT REFERRAL**

#### **A. Referring Center Responsibilities:**

1. Outpatient referral should begin with a phone call from the referring care provider to the receiving physician.
2. A convenient time for evaluation of the patient at the receiving center should be arranged.

#### **B. Receiving Center Responsibilities:**

1. The referring care provider should be contacted by either telephone or letter.
2. Whenever possible, patients should continue under the care of the referring care provider.

## **MATERNAL TRANSPORT PERSONNEL**

- I. The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and fetus.
- II. The transport team members should be selected from appropriately trained, licensed health care providers.
- III. Transport team members should have the collective expertise sufficient to provide:
  - A. monitoring of blood pressure, uterine contractions, deep tendon reflexes, and fetal heart rate
  - B. monitoring the administration of intravenous infusions and usage of tocolytic, antihypertensive, and anticonvulsant medications
  - C. care for a wide variety of emergency conditions including delivery and neonatal resuscitation.
- IV. Transport team members should be oriented to the transport vehicle and usage of transport equipment. All transport team members should follow state and national standards.
- V. In instances such as advanced labor, unstable maternal condition, or severe illness, it may become necessary for the referring physician to accompany the patient during transport, if transport is still recommended by the receiving physician. (See Appendix VI re EMTALA)

## MATERNAL TRANSPORT MODALITY

- I. Selection of the transport modality should be a joint decision by the referring and receiving physicians based on the condition of the mother and fetus.
- II. Maternal transport can be accomplished by private vehicle, ambulance, rotary wing aircraft (RWA), or fixed wing aircraft (FWA). The use of a private vehicle, RWA or FWA for maternal transport must be individualized. The ambulance (land and air) must be licensed by the Emergency Medical Services Division of the Tennessee Department of Health.
- III. If care en route is necessary, an Advanced Life Support (ALS) ambulance is required for maternal transport. The description of an ALS ambulance is defined by the Tennessee Department of Health and may be located in the latest edition of the Tennessee Emergency Medical Services Statutes and Rules, located at [www.tennessee.gov](http://www.tennessee.gov) under the "Laws & Justice" heading.

## MATERNAL TRANSPORT EQUIPMENT

- I. The referring health care provider should be aware of the availability of Advanced Life Support (ALS) ambulances in the area. The required equipment and supplies for maternal and neonatal care in an ALS ambulance are defined by the Tennessee Department of Health (See Appendix VIII).
- II. Organization and maintenance of additional transport equipment is the responsibility of the transport team.
- III. Additional equipment and supplies that may be necessary should be provided by the transporting team. These include:
  - A. doppler
  - B. reflex hammer
  - C. infusion pump
  - D. oxygen masks (premature and newborn size)
  - E. neonatal resuscitation supplies and equipment (see Appendix VII)
  - F. suction catheters (#6, #8, and #10 Fr)
  - G. latex free equipment and supplies must be available
- IV. Additional medications (or therapeutic equivalents) that may be necessary should be provided by the transport team. Such medications, including but not limited to those listed below, may be given when ordered by the referring physician.
  - A. Antenatal corticosteroids to accelerate fetal lung maturity
    1. Betamethasone for injection (12 mg IM is usual dose)
    2. Dexamethasone for injection (6 – 10 mg IM is usual dose)
  - B. Eclampsia
    1. Magnesium Sulfate 6 gram bolus IV, then 2 grams/hr IV; additional dose of 2 grams over 5-10 minutes for persistent seizures (repeat x 1)
    2. Calcium gluconate 1 gram IV to reverse magnesium overdose
    3. Sodium amobarbital 250 mg IV over 3 minutes (can substitute another short-acting benzodiazepine)
  - C. Oxytocics
    1. Oxytocin (Pitocin) 10 units per ampule / vial
    2. Misoprostol (Cytotec) 100 mcg tablets and pill cutter
    3. Methylergonovine (Methergine) 200 mcg ampules
    4. Carboprost (Hemabate) 250 microgram ampule / vial
  - D. Tocolytics
    1. Terbutaline Sulfate for injection (0.25 mg subcutaneously at 20 to 60 minute intervals is usual dose)
    2. Magnesium Sulfate (dosage same as for eclampsia)
  - E. Antibiotics for Group B strep prophylaxis (ref: CDC MMWR, August 16, 2002, / Vol.

51 / No. RR-11)

1. Penicillin G 5,000,000 units IV
2. Ampicillin 2 grams IV
3. Cefazolin 2 grams IV
4. Clindamycin 900 mg IV
5. Erythromycin 500 mg IV
6. Vancomycin 1 gram IV

F. Antihypertensives

1. Labetalol (Trandate) IV. Dosage is repeated and/or adjusted at 20 minute intervals according to patient response. May sequentially give 20 mg, then 40 mg, then 80 mg, then an additional 80 mg, if insufficient response to the lower doses. Maximum dosage is 220 mg–300 mg.
2. Hydralazine (Apresoline) 5-10 mg IV every 20 minutes. Maximum dosage is 30 mg.
3. Nifedipine (Procardia) 10 mg p.o. every 20 minutes. Maximum dosage is 100 mg.

## **MATERNAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both the referring and receiving centers have responsibilities to provide adequate documentation of clinical data.
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported patient:
    1. copy of complete prenatal record
    2. copy of current medical record (include EMTALA form)
    3. copy of record of care during transport
  - B. The referring center should maintain a record regarding disposition of transported mothers.
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported patient
  - C. send a summary of care of both mother and/or infant to the referring care provider

## EVALUATION OF PERINATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk perinatal patient requires the cooperation and coordination of many skilled health care personnel. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available through the perinatal network.
  - B. assisting perinatal care providers in developing their abilities to identify high-risk perinatal patients, anticipate complications, and stabilize those patients before transport.
  - C. continuing quality improvement through ongoing education of perinatal care and EMS providers (See Appendices I and II).
- III. Planning of the perinatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their maternal referrals with or without the assistance of the receiving center.

# **NEONATAL TRANSPORT**

## **Level II-A**

Level II-A units provide care for maternal and neonatal patients born at 34 weeks gestation or above whose courses are uncomplicated and for patients with mild obstetric and neonatal illnesses who do not require specialized services.

When antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for delivery in a facility with the needed services is encouraged.

When transport is deemed safe for the mother and fetus, transfer of mothers to a Level III unit with the specialized services required by the fetus after birth is encouraged. The elective delivery of an infant in a hospital without the required pediatric subspecialty services resulting in a planned neonatal transport is discouraged.

The nursery must have equipment and personnel to provide controlled thermal environments, hood oxygen for protracted management and assisted neonatal ventilation pending transfer to another institution for more specialized care. Obstetric and pediatric co-directors are board-certified in their respective specialties.

## **INDICATIONS FOR NEONATAL CONSULTATION AND/OR TRANSPORT**

- I.** Gestational age  $\leq$ 34 weeks
- II.** Post-asphyxial complications
- III.** Requirement for ventilatory assistance beyond the delivery room
- IV.** Requirement for total parenteral nutrition
- V.** Complicated sepsis or meningitis
- VI.** Suspected congenital heart disease
- VII.** Neurologic disorder
- VIII.** Gastrointestinal disorder
- IX.** Genitourinary disorder
- X.** Hematologic disorder
- XI.** Musculoskeletal disorder
- XII.** Endocrine or metabolic disorder
- XIII.** Congenital malformation or suspected genetic disorder requiring further evaluation

## NEONATAL REFERRAL PROCESS

I. Neonatal transport is initiated by the health care provider responsible for the patient's medical care. Guidelines for the referral process are outlined below.

### II. REFERRING CENTER RESPONSIBILITIES

- A. The referring health care provider's decision to request consultation is the first step in the referral process.
- B. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the referring health care provider in developing a treatment plan for stabilizing the patient before transport.
- C. A discussion between the referring health care provider and receiving physician regarding the neonate will result in one of three possible dispositions:
  - 1. Required neonatal care can be provided at the referring center. The receiving physician under these circumstances has only a consultative role.
  - 2. The neonate requires further observation, investigation, or other preparation before possible transport. Continued contact between the providers is necessary.
  - 3. Transport of the neonate is necessary. The optimal time, mode of transfer, transport personnel, and additional information regarding the neonate should be discussed. The process of stabilization of the neonate at the referring center should be reviewed and documented.
- D. If the neonate is transported by the referring center, the referring health care provider is responsible for the patient until arrival at the receiving center.
- E. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- F. The medical record should be organized as the patient is prepared for transport and sent with the patient.
- G. Consent forms to authorize transfer must be obtained.
- H. Parents should be encouraged to see and touch the infant. Photographs of the infant should be provided to parents, if culturally appropriate.
- I. Appropriate maternal/neonatal identification should be in place before transport.
- J. All referrals should be directly admitted to the receiving unit to avoid unnecessary delays in the emergency room.

### III. RECEIVING CENTER RESPONSIBILITIES

- A. The receiving physician is responsible for the decision to accept the referring health care provider's request for transport and make preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring health care provider in locating appropriate care.
- B. If the neonate is transported by the referring center, full responsibility begins with admission to the receiving center.
- C. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- D. If a delay in transport occurs, a repeat telephone call should be made for further assessment and advice.
- E. On arrival of the transport team, assessment and further stabilization of the neonate should be done in collaboration with the referring staff.
- F. The transport team must verify proper identification of the neonate before transport.
- G. Following stabilization of the neonate, the transport team should communicate with the parents to assure their understanding of the infant's condition, possible course, and therapeutic intervention that has been undertaken or anticipated. Information about the receiving center should be provided to the parents.
- H. Consent forms to authorize transfer, treatment, and admission to the receiving hospital must be obtained.
- I. Parents should be encouraged to see and touch the infant before departure. Photographs of the infant should be provided to parents, if culturally appropriate.
- J. Before departure, the transport team should communicate with receiving center personnel regarding the neonate's history, current status, and planned management during transport.
- K. Before arrival at the receiving center, the transport team should communicate the status of the neonate and anticipated needs on admission.
- L. A telephone call should be made to the parents shortly following admission of the neonate.
- M. Within 24 hours of admission, communication with referring center personnel regarding events during transport and since admission should occur.
- N. Periodic communication with the referring health care provider should be maintained.
- O. Consideration should be given to returning the care of the patient to the referring health care provider or primary care physician when practical and medically appropriate.
- P. Upon discharge of the infant, a discharge summary should be sent to the referring health care provider.

## NEONATAL TRANSPORT PERSONNEL

- I. The composition of the transport team should be decided jointly by the referring health care provider and receiving physician(s) based on the condition of the neonate. In most instances neonates requiring Level III care should be transported by a team meeting Level III guidelines.
- II. Transport team members should be selected from appropriately trained licensed health care providers.
- III. Transport team members should have the collective expertise sufficient to provide the following, if necessary:
  - A. observation of the neonate throughout the transport
  - B. monitoring of body temperature, respiratory status and cardiovascular status
  - C. delivery and monitoring of oxygen therapy
  - D. initiation and maintenance of IV access and therapy
  - E. positive pressure ventilation, using either a face mask or endotracheal tube
  - F. supportive care for a wide variety of emergency conditions
- IV. At least one member of the transport team should be a current Neonatal Resuscitation Program (American Academy of Pediatrics and American Heart Association) provider.
- V. Transport team members should be oriented to the transport vehicle and use of transport equipment. All transport team members should follow state and national standards.
- VI. The knowledge and skills required for a nurse to perform neonatal transport are listed in the most current edition of the Educational Objectives for Nurses, Levels I, II, III, Neonatal Transport Nurses, Tennessee Perinatal Care System, Tennessee Department of Health (Appendix I). Other suggested supplemental education includes the AAP/AHA Neonatal Resuscitation Program and The S.T.A.B.L.E. Program<sup>®</sup>, endorsed by the AAP and the March of Dimes.

## NEONATAL SPECIALTY TRANSPORT MODALITY

- I. Selection of the transport modality should be a joint decision by the referring health care provider and receiving physician based on the condition of the neonate.
- II. The ambulance used for neonatal transport must be licensed by the EMS Division of the Tennessee Department of Health at least as an ALS ambulance.
- III. The ambulance used for neonatal transport should provide:
  - A. secure fixation of the transport incubator
  - B. secure fastening of other equipment, including oxygen and air tanks
  - C. an independent power source using an inverter or generator to allow uninterrupted and fail-safe operation of the incubator and other supporting equipment
  - D. adapters required for connections to the ambulance power source.
- IV. If using a RWA or FWA, see page 73, Level III, "Neonatal Specialty Transport Modality".

## **NEONATAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both referring and receiving center personnel have responsibilities to provide adequate documentation of clinical data.
  
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported neonate:
    - 1. copy of complete maternal prenatal record
    - 2. copy of current maternal medical record
    - 3. copy of current neonatal medical record
    - 4. copies of any images in a format that can be reviewed by the transport team at the bedside
    - 5. copy of record of care during transport
  
  - B. The referring center should maintain a record regarding disposition of transported neonates.
  
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported neonate
  - C. send a summary of care to the referring perinatal health care providers

## EVALUATION OF NEONATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk neonate requires the cooperation and coordination of many skilled health care personnel. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available to them through the perinatal network.
  - B. assisting neonatal care providers in developing their abilities to identify high-risk neonatal patients, anticipate complications, and stabilize those patients before transport.
  - C. continuing quality improvement through ongoing education of neonatal care and EMS providers (See Appendices I and II).
- III. Planning of the neonatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their neonatal referrals with or without the assistance of the receiving center.

## **LEVEL II-B FACILITIES**

# **MATERNAL TRANSPORT**

## **Level II-B**

Level II-B units are capable of managing more complex maternal and neonatal abnormalities such as care of neonates that require umbilical vessel catheters and protracted mechanical ventilation. In exceptional circumstances, the Level II-B unit may receive patients transferred from Level I and Level II-A institutions.

When antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for deliver in a facility with the needed services is encouraged.

When transport is deemed safe for the mother and fetus, transfer of mothers to a Level III unit with the specialized services required by the fetus after birth is encouraged. The elective delivery of an infant in a hospital without the required pediatric subspecialty services resulting in a planned neonatal transport is discouraged.

The obstetric co-director is board certified in that specialty. The pediatric co-director is board certified in neonatal-perinatal medicine.

## INDICATIONS FOR MATERNAL CONSULTATION AND/OR TRANSPORT

### I. ANTEPARTUM

#### A. Maternal History

1. previous preterm labor (<37 weeks) or low-birthweight neonate (<2500 gm)
2. previous neonate >4000 gm at term or any large-for-gestational age neonate
3. previous stillbirth, neonatal loss, or two or more abortions
4. suspected incompetence of the cervix
5. diagnosed abnormality of the genital tract
6. medical indication for termination of previous pregnancy
7. neonate who required more than routine observation or care
8. neonate with known or suspected genetic disorder
9. severe emotional problems associated with previous pregnancy or delivery
10. previous vertical or classical uterine incision
11. age <16 or advanced maternal age ( $\geq 35$  years of age at delivery)
12. prepregnancy weight <45 kg or >90 kg
13. height <150 cm

#### B. Medical/Surgical Complications

1. diabetes mellitus/endocrine disorder
2. autoimmune disorder
3. cardiac disease
4. hypertension
5. pulmonary disease
6. renal disease
7. hematologic disorder
8. neurologic disorder
9. musculoskeletal disorder
10. infection
11. nutritional disorder
12. substance use
13. malignancy
14. psychiatric disorder
15. trauma
16. surgical emergency
17. morbid obesity

#### C. Obstetric Complications

1. glucose intolerance
2. urinary tract infection
3. sexually transmitted disease
4. positive fetal fibronectin test
5. suspected ectopic pregnancy

6. suspected missed abortion
7. hyperemesis
8. exposure to teratogen
9. isoimmunization
10. persistent anemia
11. vaginal bleeding
12. preeclampsia/eclampsia
13. suspected polyhydramnios or oligohydramnios
14. preterm cervical dilatation without uterine activity
15. preterm rupture of membranes with or without uterine activity
16. rupture of membranes at term for more than 12 hours without labor and/or evidence of amnionitis or sepsis at any time
17. suspected fetopelvic disproportion
18. inappropriate fetal growth for gestational age
19. multiple gestation
20. postterm gestation (>42 weeks)
21. fetal demise
22. known or suspected fetal anomaly
23. abnormal triple screen
24. morbid obesity

## **II. INTRAPARTUM**

- A. preterm (<37 weeks) cervical dilatation with uterine contractions
- B. abnormal presentation
- C. suspected fetopelvic disproportion
- D. dysfunctional labor
- E. rupture of membranes at term for more than 12 hours and/or evidence of amnionitis or sepsis at any time
- F. abnormal bleeding
- G. suspected nonreassuring fetal status
- H. preeclampsia/eclampsia
- I. uterine hyperstimulation syndrome
- J. meconium in amniotic fluid
- K. multiple gestation
- L. morbid obesity

## **III. POSTPARTUM**

- A. preeclampsia/eclampsia
- B. sepsis
- C. abnormal bleeding
- D. thromboembolic disease
- E. cardiopulmonary dysfunction
- F. morbid obesity
- G. neonatal transport

## **MATERNAL REFERRAL PROCESS**

I. Maternal transport is initiated by the health care provider responsible for the patient's medical care. Maternal referral may lead to admission of the patient to the receiving hospital (inpatient transport) or to outpatient evaluation and management (outpatient transport). The guidelines for the referral process are outlined below.

### **II. INPATIENT TRANSPORT**

#### **A. Referring Center Responsibilities:**

1. The decision by the referring care provider (physician, certified nurse midwife, nurse practitioner) to request consultation is the first step in the referral process.
2. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the care provider in developing a treatment plan for stabilizing the patient before and during transport.
3. Personnel of the referring and receiving facilities must follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.
4. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
5. A ground ambulance is the most appropriate vehicle for the majority of maternal transports. If an alternate form of transportation is being considered, the referring physician should discuss this alternative with the receiving physician at the time of consultation.
6. The medical record should be organized as the patient is prepared for transport and sent with the patient.
7. The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and fetus.
8. To avoid unnecessary delays in the emergency room or admitting office, all referrals should be directly admitted to the receiving obstetric unit.

#### **B. Receiving Center Responsibilities:**

1. The receiving physician is responsible for the decision to accept the referring care provider's request for transport and making preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring physician in locating appropriate alternative care.
2. Personnel of the referring and receiving facilities must follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social

Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.

3. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
4. Every patient accepted by the receiving center should be seen by a physician within 30 minutes of arrival.
5. Communication with the referring care provider should occur following admission.
6. If the patient is discharged undelivered, communication should occur prior to the time of discharge.
7. A summary of care of both mother and infant should be sent to the referring care provider.
8. Every effort should be made to return the patient to the care of the referring care provider as soon as possible.

### **III. OUTPATIENT REFERRAL**

#### **A. Referring Center Responsibilities:**

1. Outpatient referral should begin with a phone call from the referring care provider to the receiving physician.
2. A convenient time for evaluation of the patient at the receiving center should be arranged.

#### **B. Receiving Center Responsibilities:**

1. The referring care provider should be contacted by either telephone or letter.
2. Whenever possible, patients should continue under the care of the referring care provider.

## **MATERNAL TRANSPORT PERSONNEL**

- I. The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and fetus.
- II. The transport team members should be selected from appropriately trained, licensed health care providers.
- III. Transport team members should have the collective expertise sufficient to provide:
  - A. monitoring of blood pressure, uterine contractions, deep tendon reflexes, and fetal heart rate
  - B. monitoring the administration of intravenous infusions and usage of tocolytic, antihypertensive, and anticonvulsant medications
  - C. care for a wide variety of emergency conditions including delivery and neonatal resuscitation.
- IV. Transport team members should be oriented to the transport vehicle and usage of transport equipment. All transport team members should follow state and national standards.
- V. In instances such as advanced labor, unstable maternal condition, or severe illness, it may become necessary for the referring physician to accompany the patient during transport, if transport is still recommended by the receiving physician.

## MATERNAL TRANSPORT MODALITY

- I. Selection of the transport modality should be a joint decision by the referring and receiving physicians based on the condition of the mother and fetus.
- II. Maternal transport can be accomplished by private vehicle, ambulance, rotary wing aircraft (RWA), or fixed wing aircraft (FWA). The use of a private vehicle, RWA or FWA for maternal transport must be individualized. The ambulance (land and air) must be licensed by the Emergency Medical Services Division of the Tennessee Department of Health.
- III. If care en route is necessary, an Advanced Life Support (ALS) ambulance is required for maternal transport. The description of an ALS ambulance is defined by the Tennessee Department of Health and may be located in the Tennessee Emergency Medical Services Statutes and Rules, latest edition, located at [www.tennessee.gov](http://www.tennessee.gov) under the "Laws & Justice" heading.

## MATERNAL TRANSPORT EQUIPMENT

- I. The referring health care provider should be aware of the availability of Advanced Life Support (ALS) ambulances in the area. The required equipment and supplies for maternal and neonatal care in an ALS ambulance are defined by the Tennessee Department of Health (See Appendix VIII.).
- II. Organization and maintenance of additional transport equipment is the responsibility of the transport team.
- III. Additional equipment and supplies that may be necessary should be provided by the transporting team. These include:
  - A. doppler
  - B. reflex hammer
  - C. infusion pump
  - D. oxygen masks (premature and newborn size)
  - E. neonatal resuscitation supplies and equipment (see Appendix VII)
  - F. suction catheters (#6, #8, and #10 Fr)
  - G. latex free equipment and supplies must be available
- IV. Additional medications (or therapeutic equivalents) that may be necessary should be provided by the transport team. Such medications, including but not limited to those listed below, may be given when ordered by the referring physician.
  - A. Antenatal corticosteroids to accelerate fetal lung maturity
    1. Betamethasone for injection (12 mg IM is usual dose)
    2. Dexamethasone for injection (6–10 mg IM is usual dose)
  - B. Eclampsia
    1. Magnesium Sulfate 6 gram bolus IV, then 2 grams/hr IV; additional dose of 2 grams over 5-10 minutes for persistent seizures (repeat x 1)
    2. Calcium gluconate 1 gram IV to reverse magnesium overdose
    3. Sodium amobarbital 250 mg IV over 3 minutes (can substitute another short-acting benzodiazepine)
  - C. Oxytocics
    1. Oxytocin (Pitocin) 10 units per ampule / vial
    2. Misoprostol (Cytotec) 100 mcg tablets and pill cutter
    3. Methylergonovine (Methergine) 200 mcg ampules
    4. Carboprost (Hemabate) 250 microgram ampule / vial
  - D. Tocolytics
    1. Terbutaline Sulfate for injection (0.25 mg subcutaneously at 20 to 60 minute intervals is usual dose)
    2. Magnesium Sulfate (dosage same as for eclampsia)
  - E. Antibiotics for Group B strep prophylaxis (ref: CDC MMWR, August 16, 2002, / Vol.

51 / No. RR-11)

1. Penicillin G 5,000,000 units IV
2. Ampicillin 2 grams IV
3. Cefazolin 2 grams IV
4. Clindamycin 900 mg IV
5. Erythromycin 500 mg IV
6. Vancomycin 1 gram IV

F. Antihypertensives

1. Labetalol (Trandate) IV. Dosage is repeated and/or adjusted at 20 minute intervals according to patient response. May sequentially give 20 mg, then 40 mg, then 80 mg, then an additional 80 mg, if insufficient response to the lower doses. Maximum dosage is 220 mg–300 mg.
2. Hydralazine (Apresoline) 5-10 mg IV every 20 minutes. Maximum dosage is 30 mg.
3. Nifedipine (Procardia) 10 mg p.o. every 20 minutes. Maximum dosage is 100 mg.

## **MATERNAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both the referring and receiving centers have responsibilities to provide adequate documentation of clinical data.
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported patient:
    - 1. copy of complete prenatal record
    - 2. copy of current medical record (include EMTALA form)
    - 3. copy of record of care during transport
  - B. The referring center should maintain a record regarding disposition of transported mothers.
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported patient
  - C. send a summary of care of both mother and/or infant to the referring care provider

## EVALUATION OF PERINATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk perinatal patient requires the cooperation and coordination of many skilled health care personnel. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available through the perinatal network.
  - B. assisting perinatal care providers in developing their abilities to identify high-risk perinatal patients, anticipate complications, and stabilize those patients before transport.
  - C. continuing quality improvement through ongoing education of perinatal care and EMS providers (See Appendices I and II).
- III. Planning of the perinatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their maternal referrals with or without the assistance of the receiving center.

# **NEONATAL TRANSPORT**

## **Level II-B**

Level II-B units are capable of managing more complex maternal and neonatal abnormalities such as care of neonates that require umbilical vessel catheters and protracted mechanical ventilation. In exceptional circumstances, the Level II-B unit may receive patients transferred from Level I and Level II-A institutions.

When antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for deliver in a facility with the needed services is encouraged.

When transport is deemed safe for the mother and fetus, transfer of mothers to a Level III unit with the specialized services required by the fetus after birth is encouraged. The elective delivery of an infant in a hospital without the required pediatric sub-specialty services resulting in a planned neonatal transport is discouraged.

The obstetric co-director is board certified in that specialty. The pediatric co-director is board certified in neonatal-perinatal medicine.

## **INDICATIONS FOR NEONATAL CONSULTATION AND/OR TRANSPORT**

- I.** Gestational age  $\leq$  28 weeks
- II.** Should be considered if birth weight is  $<1000$  grams and must occur if birthweight is  $<750$  gm
- III.** Severity of illness requiring a level of care that exceeds the capacity of the Level II-B facility

## NEONATAL REFERRAL PROCESS

- I. Neonatal transport is initiated by the health care provider responsible for the patient's medical care. Guidelines for the referral process are outlined below.

### II. REFERRING CENTER RESPONSIBILITIES

- A. The referring health care provider's decision to request consultation is the first step in the referral process.
- B. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the referring health care provider in developing a treatment plan for stabilizing the patient before transport.
- C. A discussion between the referring health care provider and receiving physician regarding the neonate will result in one of three possible dispositions:
  1. Required neonatal care can be provided at the referring center. The receiving physician under these circumstances has only a consultative role.
  2. The neonate requires further observation, investigation, or other preparation before possible transport. Continued contact between the providers is necessary.
  3. Transport of the neonate is necessary. The optimal time, mode of transfer, transport personnel, and additional information regarding the neonate should be discussed. The process of stabilization of the neonate at the referring center should be reviewed and documented.
- D. If the neonate is transported by the referring center, the referring health care provider is responsible for the patient until arrival at the receiving center.
- E. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- F. The medical record should be organized as the patient is prepared for transport and sent with the patient.
- G. Consent forms to authorize transfer must be obtained.
- H. Parents should be encouraged to see and touch the infant. Photographs of the infant should be provided to parents, if culturally appropriate.
- I. Appropriate maternal/neonatal identification should be in place before transport.
- J. All referrals should be directly admitted to the receiving unit to avoid unnecessary delays in the emergency room.

### III. RECEIVING CENTER RESPONSIBILITIES

- A. The receiving physician is responsible for the decision to accept the referring health care provider's request for transport and make preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring health care provider in locating appropriate care.
- B. If the neonate is transported by the referring center, full responsibility begins with admission to the receiving center.
- C. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- D. If a delay in transport occurs, a repeat telephone call should be made for further assessment and advice.
- E. On arrival of the transport team, assessment and further stabilization of the neonate should be done in collaboration with the referring staff.
- F. The transport team must verify proper identification of the neonate before transport.
- G. Following stabilization of the neonate, the transport team should communicate with the parents to assure their understanding of the infant's condition, possible course, and therapeutic intervention that has been undertaken or anticipated. Information about the receiving center should be provided to the parents.
- H. Consent forms to authorize transfer, treatment, and admission to the receiving hospital must be obtained.
- I. Parents should be encouraged to see and touch the infant before departure. Photographs of the infant should be provided to parents, if culturally appropriate.
- J. Before departure, the transport team should communicate with receiving center personnel regarding the neonate's history, current status, and planned management during transport.
- K. Before arrival at the receiving center, the transport team should communicate the status of the neonate and anticipated needs on admission.
- L. A telephone call should be made to the parents shortly following admission of the neonate.
- M. Within 24 hours of admission, communication with referring center personnel regarding events during transport and since admission should occur.
- N. Periodic communication with the referring health care provider should be maintained.
- O. Consideration should be given to returning the care of the patient to the referring health care provider or primary care physician when practical and medically appropriate.
- P. Upon discharge of the infant, a discharge summary should be sent to the referring health care provider.

## NEONATAL TRANSPORT PERSONNEL

- I. The composition of the transport team should be decided jointly by the referring health care provider and receiving physician(s) based on the condition of the neonate.
- II. Transport team members should be selected from appropriately trained physicians, neonatal nurse practitioners, registered nurses, respiratory therapists, emergency medical technicians, and paramedics. The minimum number of transport team members should be three, of which one is the operator of the transport vehicle. The other team members should be comprised of a neonatologist, neonatal fellow, neonatal nurse practitioner, neonatal specialty trained registered nurse, or neonatal specialty trained respiratory therapist.
- III. Transport team members should have the collective expertise sufficient to provide, if necessary:
  - A. observation of the neonate throughout the transport
  - B. monitoring of body temperature, respiratory status, and cardiovascular status
  - C. delivery and monitoring of oxygen therapy
  - D. advanced airway management and mechanical ventilation
  - E. advanced intravascular access and management skills
  - F. recognition and management of a wide variety of emergency conditions
  - G. administration of rescue and stabilization medication
- IV. At least two members of the transport team should be current Neonatal Resuscitation Program (American Academy of Pediatrics and American Heart Association) providers. It is also recommended that team members maintain current S.T.A.B.L.E. training.
- V. Transport team members should be oriented to the transport vehicle and use of transport equipment. All transport team members should follow state and national standards.
- VI. The knowledge and skills required for a nurse to perform neonatal transport are listed in the most current edition of Educational Objectives for Nurses, Levels I, II, III Neonatal Transport Nurses, Tennessee Perinatal Care System, Tennessee Department of Health (Appendix I).

## NEONATAL SPECIALTY TRANSPORT MODALITY

- I. Selection of the mobile neonatal intensive care unit should be based on the condition of the neonate.
- II. The vehicle used exclusively for the provision of neonatal intensive care and transportation between medical facilities shall conform with the following standards for design and construction as defined by the EMS Division of the Tennessee Department of Health (Tennessee Emergency Medical Services Statutes and Rules, Rule 1200-12-1-.02[4] Special Vehicle Requirements).
  - A. Exterior surfaces, emblems, and markings shall conform to Federal Specifications-Ambulance.
  - B. Additional markings, legends, or logos may be used to identify the provider and purpose for specific vehicles, except that no letter shall exceed 14 inches in height. Legends as "neonatal intensive care or critical care transport" may be substituted for the "Ambulance" in exterior markings.
  - C. Warning lights and siren shall be furnished in accordance with Federal Specification-Ambulance except that side floodlights shall not be required.
  - D. Vehicle crashworthiness shall be assured with roll-cage construction evidenced by compliance with the Ambulance Manufacturer's Division Standards of the Truck Body and Equipment Association or comparable construction under written statement and performance bond by the manufacturer.
  - E. Doors shall provide access to the rear and curbside of the patient compartment. Where the vertical lift distance of the patient loading area exceeds 28 inches, a ramp or electrical-hydraulic lift shall be furnished to facilitate patient loading.
  - F. Environmental systems on the unit shall meet heating/air-conditioning standards as specified in Federal Specifications-Ambulance.
  - G. Vehicle electrical systems shall be provided to furnish 110 volt AC power sufficient to sustain 3,000 watts at 60 cycles. The unit shall be equipped with a back-up power system sufficient to operate patient care equipment in the event of failure of the main power systems. The 110 volt system shall incorporate a ground fault interrupter device for protection against electrical hazards.
  - H. Patient compartment shall be so designed to provide the following:
    1. a properly secured transport incubator allowing observation from at least two sides
    2. compartments for appropriate storage of materials
    3. illumination at the primary patient care area of at least 75 foot candles
    4. safety features such as sculpted, padded, or recessed cabinet corners and latches to prevent undue injury during sudden deceleration
    5. safety devices such as a grab rail or hand strap, secured according to Federal Motor Vehicle Safety Standards for safety restraints, safety belts provided at all attendant seats, and safety restraint devices for neonates
  - I. An oxygen system with sufficient capacity to deliver a minimum continuous flow of 10 liters per minute for at least four hours must be available. The installed oxygen system shall be capable of delivering specific monitored blended oxygen concentrations.
  - J. The vehicle shall provide environmental conditions for the neonate that minimize the risk of temperature instability and excessive noise and vibration.

- III. In the event a RWA or FWA is used for neonatal transport, the following additional precautions should be observed:
  - A. use of a licensed air medical transport program with appropriately credentialed neonatal care providers
  - B. secure fastening of transport equipment
  - C. an independent power source to allow uninterrupted and fail-safe operation of the incubator and other supporting equipment
  - D. environmental conditions for the neonate that minimize the risk of temperature instability, noise and vibration
  - E. protective headgear for crew use during transport
  - F. system for easy communication among the members of the transport team and medical control at receiving center
  - G. seat belts
  - H. seating arrangement that permits close observation and handling of the patient
- IV. Latex free equipment and supplies must be provided.

## NEONATAL TRANSPORT EQUIPMENT

- I. Organization and maintenance of neonatal transport equipment is the responsibility of the transporting facility. (See Appendix VIII)
- II. Equipment to maintain a neutral thermal environment for the neonate should include:
  - A. transport incubator
  - B. thermometer
  - C. blanket, insulating blanket or chemically activated heat pack (appropriate for neonatal use)
- III. The transport incubator should meet the following requirements:
  - A. approved by the manufacturer for use during transport and installed in the transport vehicle with crashworthy restraints.
  - B. if used in RWA and FWA it shall meet FAA requirements for crashworthiness and flammability of materials. The transport incubator and monitoring equipment should be tested by an FAA certified mechanic to assure equipment does not interfere with navigational instruments.
  - C. a heat source that requires minimal time for preheating and should maintain ambient temperature within the desired range of 29° to 36° C. The control for temperature setting should be readily accessible and easy to operate, and there should be provision for easy determination of ambient temperature. It is essential to have a fail-safe alarm system that will recognize overheating or underheating.
  - D. provide an environment in which the oxygen supply is constant and controllable.
  - E. provide unrestricted visibility of the neonate with a functional independent light source for general illumination provided in or on the incubator.
  - F. easy accessibility to the neonate resulting in minimal interference with thermal protection and oxygen supply.
  - G. safety restraint devices to secure the neonate inside the incubator that conform to Federal Motor Vehicle Safety Standards and FAA Standards.
- IV. Equipment for oxygen delivery and monitoring should include:
  - A. oxygen tanks
  - B. air tanks
  - C. pressure gauges
  - D. flowmeters (needle control valve preferred)
  - E. oxygen analyzer
  - F. oxygen blender
  - G. oxygen tubing and adapters
  - H. oxygen hood or nasal cannula
  - I. neonatal oxygen mask
  - J. neonatal resuscitation manual bag and mask with manometer or T-piece resuscitator
  - K. continuous positive airway pressure apparatus
  - L. mechanical ventilator
- V. The following guidelines are related to the use of oxygen during transport:

- A. a portable supply of oxygen and compressed air in cylinders adequate to last the entire journey with surplus to cover unexpected needs and delays should be carried. Proper restraint of these cylinders is mandatory throughout the transport. An oxygen cylinder usage chart may be used in determining the number and type of oxygen cylinders necessary during transport (Appendix III).
  - B. oxygen cylinders in use should be provided with pressure gauges and flow meters.
  - C. the ambient oxygen concentration must be monitored continuously by an oxygen analyzer.
- VI.** Devices to maintain the patency of the airway and gastric decompression must be readily available and should include:
- A. bulb syringe
  - B. regulated suction with gauge
  - C. suction catheters (#6, 8, 10 Fr)
  - D. feeding tube (#8 Fr) with a 20 mL syringe
  - E. Replogle tube (#8 Fr)
- VII.** Equipment for vital sign monitoring should include:
- A. a continuous heart rate monitor
  - B. neonatal stethoscope
  - C. body temperature monitor
  - D. noninvasive and invasive blood pressure monitoring devices
  - E. pulse oximeter for noninvasive monitoring of oxygen saturation
- VIII.** Equipment for monitoring blood glucose must be available.
- IX.** If intravenous therapy is required, an infusion pump that is portable, battery-powered, fail-safe, and calibrated to ensure accurate delivery of calculated fluid volumes must be used. Critical neonates may require multiple intravenous lines requiring additional infusion pumps.
- X.** The equipment and supplies required for resuscitation of a neonate must be available, portable and should include:
- A. Endotracheal intubation
    - 1. laryngoscope handle with blades (#00, 0, 1)
    - 2. laryngoscope spare bulbs
    - 3. laryngoscope spare batteries
    - 4. endotracheal tubes (#2.5, 3.0, 3.5, 4.0 mmID)
    - 5. neonatal resuscitation manual bag and mask or T-piece resuscitator
    - 6. pressure manometer (to monitor PIP and PEEP if manual ventilation is necessary)
    - 7. disposable stylet (#6 Fr)
    - 8. adhesive tape or commercial endotracheal tube holders
    - 9. scissors
    - 10. end tidal carbon dioxide (CO<sub>2</sub>) detector
  - B. Intravenous infusion

1. intravenous needles and catheters (#22, 23, 24, 25, 26 gauge)
  2. syringes (1, 3, 6, 12, 20, 35 mL)
  3. intravenous armboard
  4. intravenous tubing and T connector
  5. infusion device
  6. tape or site dressing
  7. site preps
  8. 3-way stopcock
  9. umbilical catheterization equipment
- C. Medications should include the following drugs or an approved therapeutic equivalent based on local program protocols.
1. adenosine
  2. ampicillin
  3. atropine
  4. calcium gluconate
  5. dextrose solution (D5W)
  6. dextrose solution D10W)
  7. dextrose solution (D25W)
  8. dobutamine
  9. dopamine
  10. epinephrine (1:10,000)
  11. fentanyl
  12. furosemide
  13. gentamicin
  14. glucagon
  15. heparin
  16. insulin (regular)
  17. morphine sulfate
  18. naloxone hydrochloride
  19. normal saline
  20. phenobarbital
  21. prostaglandin E<sub>1</sub> (requires refrigeration)
  22. sodium bicarbonate (4.2%)
  23. sterile water
  24. surfactant (requires refrigeration)
  25. THAM or other buffer

**XI. Equipment for diagnosis and management of air leak syndrome**

- A. transilluminator
- B. chest tubes (8, 10, 12 Fr)
- C. chest tube kit
- D. 18 to 20 gauge over the needle catheters
- E. drain with one-way check valve

- F. 3-way stopcock
- G. luer lock and slip tip syringe (35/60cc)

**XII.** Equipment for handwashing and personal protection

- A. antiseptic solution or towelettes
- B. gloves
- C. full face protection or goggles and masks
- D. fluid-retardant and fluid-resistant gowns
- E. infectious waste disposal bags
- F. soiled linen disposal bags
- G. sharps box

**XIII.** Latex free equipment and supplies must be provided.

## **NEONATAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both referring and receiving center personnel have responsibilities to provide adequate documentation of clinical data.
  
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported neonate:
    - 1. copy of complete maternal prenatal record
    - 2. copy of current maternal medical record
    - 3. copy of current neonatal medical record
    - 4. copies of any images in a format that can be reviewed by the transport team at the bedside
    - 5. copy of record of care during transport
  
  - B. The referring center should maintain a record regarding disposition of transported neonates.
  
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported neonate
  - C. send a summary of care to the referring perinatal health care providers

## EVALUATION OF NEONATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk neonate requires the cooperation and coordination of many skilled persons. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available to them through the perinatal network.
  - B. assisting neonatal care providers in developing their abilities to identify high-risk neonatal patients, anticipate complications, and stabilize those patients before transport.
  - C. continuing quality improvement through ongoing education of neonatal care and EMS providers (See Appendices I and II).
- III. Planning of the neonatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their neonatal referrals.

## **LEVEL III FACILITIES**

# **MATERNAL TRANSPORT**

## **Level III**

Level III units have the capacity to manage the most complex and severe maternal and neonatal illnesses by virtue of their equipment, perinatal staff and on-site availability of a complete spectrum of pediatric sub-specialists. Consultation with specialized physicians elsewhere should rarely be necessary.

Although it is not always possible to prenatally anticipate the need for pediatric subspecialty services, when antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for delivery in a facility with the needed services is encouraged.

The elective and/or planned delivery of a fetus with a condition(s) that require immediate neonatal transport should be discouraged.

The Level III unit, if it so chooses, is responsible for providing equipment and qualified staff to transport sick infants from other referring hospitals.

The Level III facility is responsible for delivery of a formal ongoing program of education in obstetrics and neonatal-perinatal medicine for its staff.

The obstetric co-director must be board-certified in maternal-fetal medicine. The pediatric co-director must be board-certified in neonatal-perinatal medicine.

### **REGIONAL PERINATAL CENTERS**

Each of Tennessee's five Regional Perinatal Centers (Northeast Tennessee Regional Perinatal Center in Johnson City; East Tennessee Regional Perinatal Center in Knoxville; Southeast Regional Perinatal Center in Chattanooga; Middle Tennessee Regional Perinatal Center in Nashville; and West Tennessee Regional Perinatal Center in Memphis) is capable of providing Level III obstetric and neonatal care.

In addition, each Regional Perinatal Center must provide the services of consultation/referral, professional education, maternal-fetal and neonatal transport, site visits upon request, post-neonatal follow-up, and data collection.

## **INDICATIONS FOR MATERNAL CONSULTATION AND/OR TRANSPORT**

- I. Maternal or fetal conditions requiring or potentially requiring specific medical or surgical services unavailable at the Level III facility. Such consultation/transfer may necessitate out of state consultation/transfer.

## MATERNAL REFERRAL PROCESS

I. Maternal transport is initiated by the health care provider responsible for the patient's medical care. Maternal referral may lead to admission of the patient to the receiving hospital (inpatient transport) or to outpatient evaluation and management (outpatient transport). The guidelines for the referral process are outlined below.

### II. INPATIENT TRANSPORT

#### A. Referring Center Responsibilities:

1. The decision by the referring care provider (physician, certified nurse midwife, nurse practitioner) to request consultation is the first step in the referral process.
2. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the care provider in developing a treatment plan for stabilizing the patient before and during transport.
3. Personnel of the referring and receiving facilities follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.
4. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
5. A ground ambulance is the most appropriate vehicle for the majority of maternal transports. If an alternate form of transportation is being considered, the referring physician should discuss this alternative with the receiving physician at the time of consultation.
6. The medical record should be organized as the patient is prepared for transport and sent with the patient.
7. The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and fetus.
8. To avoid unnecessary delays in the emergency room or admitting office, all referrals should be directly admitted to the receiving obstetric unit.

#### B. Receiving Center Responsibilities:

1. The receiving physician is responsible for the decision to accept the referring care provider's request for transport and making preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring physician in locating appropriate alternative care.
2. Personnel of the referring and receiving facilities follow the COBRA/EMTALA guidelines (42USC 1395dd. Section 1867 of the Social Security Act. Also known as Section 9121 of the Consolidated Omnibus Budget Reconciliation

Act of 1985. See Appendix VI). Regional Perinatal Centers shall not refuse to accept an appropriate transfer of an individual who requires such specialized care if the hospital has the capability and bed availability to treat the individual. All transfers should be based on medical need.

3. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics).
4. Every patient accepted by the receiving center should be seen by a physician within 30 minutes of arrival.
5. Communication with the referring care provider should occur following admission.
6. If the patient is discharged undelivered, communication should occur prior to the time of discharge.
7. A summary of care of both mother and infant should be sent to the referring care provider.
8. Every effort should be made to return the patient to the care of the referring care provider as soon as possible.

### **III. OUTPATIENT REFERRAL**

#### **A. Referring Center Responsibilities:**

1. Outpatient referral should begin with a phone call from the referring care provider to the receiving physician.
2. A convenient time for evaluation of the patient at the receiving center should be arranged.

#### **B. Receiving Center Responsibilities:**

1. The referring care provider should be contacted by either telephone or letter.
2. Whenever possible, patients should continue under the care of the referring care provider.

## MATERNAL TRANSPORT PERSONNEL

- I. The composition of the transport team should be decided jointly by the referring and receiving care providers based on the condition of the mother and fetus.
- II. The transport team members should be selected from appropriately trained, licensed health care providers.
- III. Transport team members should have the collective expertise sufficient to provide the following, if necessary:
  - A. monitoring of blood pressure, uterine contractions, deep tendon reflexes, and fetal heart rate
  - B. monitoring the administration of intravenous infusions and usage of tocolytic, antihypertensive, and anticonvulsant medications
  - C. care for a wide variety of emergency conditions including delivery and neonatal resuscitation.
- IV. Transport team members should be oriented to the transport vehicle and usage of transport equipment. All transport team members should follow state and national standards.
- V. In instances such as advanced labor, unstable maternal condition, or severe illness, it may become necessary for the referring physician to accompany the patient during transport, if transport is still recommended by the receiving physician.

## MATERNAL TRANSPORT MODALITY

- I. Selection of the transport modality should be a joint decision by the referring and receiving physicians based on the condition of the mother and fetus.
- II. Maternal transport can be accomplished by private vehicle, ambulance, rotary wing aircraft (RWA), or fixed wing aircraft (FWA). The use of a private vehicle, RWA or FWA for maternal transport must be individualized. The ambulance (land and air) must be licensed by the Emergency Medical Services Division of the Tennessee Department of Health.
- III. If care en route is necessary, an Advanced Life Support (ALS) ambulance is required for maternal transport. The description of an ALS ambulance is defined by the Tennessee Department of Health and may be located in the latest edition of the Tennessee Emergency Medical Services Statutes and Rules, located at [www.tennessee.gov](http://www.tennessee.gov) under the "Laws & Justice" heading.

## MATERNAL TRANSPORT EQUIPMENT

- I. The referring health care provider should be aware of the availability of Advanced Life Support (ALS) ambulances in the area. The required equipment and supplies for maternal and neonatal care in an ALS ambulance are defined by the Tennessee Department of Health (See Appendix VIII).
- II. Organization and maintenance of additional transport equipment is the responsibility of the transport team.
- III. Additional equipment and supplies that may be necessary should be provided by the transporting team. These include:
  - A. doppler
  - B. reflex hammer
  - C. infusion pump
  - D. oxygen masks (premature and newborn size)
  - E. neonatal resuscitation supplies and equipment (see Appendix VII)
  - F. suction catheters (#6, #8, and #10 Fr)
  - G. latex free equipment and supplies must be available, when possible.
- IV. Additional medications (or therapeutic equivalents) that may be necessary should be provided by the transport team. Such medications, including but not limited to those listed below, may be given when ordered by the referring physician.
  - A. Antenatal corticosteroids to accelerate fetal lung maturity
    1. Betamethasone for injection (12 mg IM is usual dose)
    2. Dexamethasone for injection (6–10 mg IM is usual dose)
  - B. Eclampsia
    1. Magnesium Sulfate 6 gram bolus IV, then 2 grams/hr IV; additional dose of 2 grams over 5-10 minutes for persistent seizures (repeat x 1)
    2. Calcium gluconate 1 gram IV to reverse magnesium overdose
    3. Sodium amobarbital 250 mg IV over 3 minutes (can substitute another short-acting benzodiazepine)
  - C. Oxytocics
    1. Oxytocin (Pitocin) 10 units per ampule / vial
    2. Misoprostol (Cytotec) 100 mcg tablets and pill cutter
    3. Methylergonovine (Methergine) 200 mcg ampules
    4. Carboprost (Hemabate) 250 microgram ampule / vial
  - D. Tocolytics
    1. Terbutaline Sulfate for injection (0.25 mg subcutaneously at 20 to 60 minute intervals is usual dose)
    2. Magnesium Sulfate (dosage same as for eclampsia)
  - E. Antibiotics for Group B strep prophylaxis (ref: CDC MMWR, August 16, 2002, / Vol.

51 / No. RR-11)

1. Penicillin G 5,000,000 units IV
2. Ampicillin 2 grams IV
3. Cefazolin 2 grams IV
4. Clindamycin 900 mg IV
5. Erythromycin 500 mg IV
6. Vancomycin 1 gram IV

F. Antihypertensives

1. Labetalol (Trandate) IV. Dosage is repeated and/or adjusted at 20 minute intervals according to patient response. May sequentially give 20 mg, then 40 mg, then 80 mg, then an additional 80 mg, if insufficient response to the lower doses. Maximum dosage is 220 mg–300 mg.
2. Hydralazine (Apresoline) 5-10 mg IV every 20 minutes. Maximum dosage is 30 mg.
3. Nifedipine (Procardia) 10 mg p.o. every 20 minutes. Maximum dosage is 100 mg.

## **MATERNAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both the referring and receiving centers have responsibilities to provide adequate documentation of clinical data.
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported patient:
    1. copy of complete prenatal record
    2. copy of current medical record (include EMTALA form)
    3. copy of record of care during transport
  - B. The referring center should maintain a record regarding disposition of transported mothers.
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported patient
  - C. send a summary of care of both mother and/or infant to the referring care provider

## EVALUATION OF PERINATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk perinatal patient requires the cooperation and coordination of many skilled health care personnel. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available through the perinatal network
  - B. assisting perinatal care providers in developing their abilities to identify high-risk perinatal patients, anticipate complications, and stabilize those patients before transport
  - C. continuing quality improvement through ongoing education of perinatal care and EMS providers (See Appendices I and II).
- III. Planning of the perinatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their maternal referrals with or without the assistance of the receiving center.

# **NEONATAL TRANSPORT**

## **Level III**

Level III units have the capacity to manage the most complex and severe maternal and neonatal illnesses by virtue of their equipment, perinatal staff and on-site availability of a complete spectrum of pediatric sub-specialists. Consultation with specialized physicians elsewhere should rarely be necessary.

Although it is not always possible to prenatally anticipate the need for pediatric subspecialty services, when antenatal ultrasound and/or genetic testing has identified a fetus with congenital anomalies, prenatal referral to an appropriate subspecialty provider or fetal assessment clinic to provide families with prognostic information and facilitate a coordinated plan for delivery in a facility with the needed services is encouraged.

The elective and/or planned delivery of a fetus with a condition(s) that require immediate neonatal transport should be discouraged.

The Level III unit, if it so chooses, is responsible for providing equipment and qualified staff to transport sick infants from other referring hospitals.

The Level III facility is responsible for delivery of a formal ongoing program of education in obstetrics and neonatal-perinatal medicine for its staff.

The obstetric co-director must be board-certified in maternal-fetal medicine. The pediatric co-director must be board-certified in neonatal-perinatal medicine.

### **REGIONAL PERINATAL CENTERS**

Each of Tennessee's five Regional Perinatal Centers (Northeast Tennessee Regional Perinatal Center in Johnson City; East Tennessee Regional Perinatal Center in Knoxville; Southeast Tennessee Regional Perinatal Center in Chattanooga; Middle Tennessee Regional Perinatal Center in Nashville; and West Tennessee Regional Perinatal Center in Memphis) is capable of providing Level III obstetric and neonatal care.

In addition, each Regional Perinatal Center must provide the services of consultation/referral; professional education; maternal-fetal and neonatal transport; site visits upon request; post-neonatal follow-up; and data collection.

## **INDICATIONS FOR NEONATAL CONSULTATION AND/OR TRANSPORT**

- I. Neonatal conditions requiring or potentially requiring specific medical or surgical services unavailable at the Level III facility. Such consultation/transfer may necessitate out of state consultation/transfer.

## NEONATAL REFERRAL PROCESS

- I. Neonatal transport is initiated by the health care provider responsible for the patient's medical care. Guidelines for the referral process are outlined below.

### II. REFERRING CENTER RESPONSIBILITIES

- A. The referring health care provider's decision to request consultation is the first step in the referral process.
- B. Telephone consultation with the receiving physician is necessary to initiate the referral process and to prepare the receiving center. This consultation may aid the referring health care provider in developing a treatment plan for stabilizing the patient before transport.
- C. A discussion between the referring health care provider and receiving physician regarding the neonate will result in one of three possible dispositions.
  1. Required neonatal care can be provided at the referring center. The receiving physician under these circumstances has only a consultative role.
  2. The neonate requires further observation, investigation, or other preparation before possible transport. Continued contact between the providers is necessary.
  3. Transport of the neonate is necessary. The optimal time, mode of transfer, transport personnel, and additional information regarding the neonate should be discussed. The process of stabilization of the neonate at the referring center should be reviewed and documented.
- D. If the neonate is transported by the referring center, the referring health care provider is responsible for the patient until arrival at the receiving center.
- E. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- F. The medical record should be organized as the patient is prepared for transport and sent with the patient.
- G. Consent forms to authorize transfer must be obtained.
- H. Parents should be encouraged to see and touch the infant. Photographs of the infant should be provided to parents, if culturally appropriate.
- I. Appropriate maternal/neonatal identification should be in place before transport.
- J. All referrals should be directly admitted to the receiving unit to avoid unnecessary delays in the emergency room.

### III. RECEIVING CENTER RESPONSIBILITIES

- A. The receiving physician is responsible for the decision to accept the referring health care provider's request for transport and make preparations at the receiving center. If unable to accept the transport, assistance will be provided to the referring health care provider in locating appropriate care.
- B. If the neonate is transported by the referring center, full responsibility begins with admission to the receiving center.
- C. If the neonate is transported by the receiving center, the referring health care provider is responsible for the patient until the arrival of the transport team. If the transport team is sent by the receiving hospital, the receiving physician or designee assumes responsibility for patient care from the time the patient leaves the referring hospital. It should be emphasized that during the preparation for transport by the transport team, the referring physician and hospital have retained responsibility for the patient unless there have been other prior agreements which determine this responsibility (most recent edition, Guidelines for Perinatal Care, published by the American College of Obstetricians and Gynecologists and the American Academy of Pediatrics). The patient becomes the full responsibility of the transport team when leaving the referring center.
- D. If a delay in transport occurs, a repeat telephone call should be made for further assessment and advice.
- E. On arrival of the transport team, assessment and further stabilization of the neonate should be done in collaboration with the referring staff.
- F. The transport team must verify proper identification of the neonate before transport.
- G. Following stabilization of the neonate, the transport team should communicate with the parents to assure their understanding of the infant's condition, possible course, and therapeutic intervention that has been undertaken or anticipated. Information about the receiving center should be provided to the parents.
- H. Consent forms to authorize transfer, treatment, and admission to the receiving hospital must be obtained.
- I. Parents should be encouraged to see and touch the infant before departure. Photographs of the infant should be provided to parents, if culturally appropriate.
- J. Before departure, the transport team should communicate with receiving center personnel regarding the neonate's history, current status, and planned management during transport.
- K. Before arrival at the receiving center, the transport team should communicate the status of the neonate and anticipated needs on admission.
- L. A telephone call should be made to the parents shortly following admission of the neonate.
- M. Within 24 hours of admission, communication with referring center personnel regarding events during transport and since admission should occur.
- N. Periodic communication with the referring health care provider should be maintained.
- O. Consideration should be given to returning the care of the patient to the referring health care provider or primary care physician when practical and medically appropriate.
- P. Upon discharge of the infant, a discharge summary should be sent to the referring health care provider.

## NEONATAL TRANSPORT PERSONNEL

- I. The composition of the transport team should be decided jointly by the referring health care provider and receiving physician(s) based on the condition of the neonate.
- II. Transport team members should be selected from appropriately trained physicians, neonatal nurse practitioners, registered nurses, respiratory therapists, emergency medical technicians, and paramedics. The minimum number of transport team members should be three, of which one is the operator of the transport vehicle. The other team members should be comprised of a neonatologist, neonatal fellow, neonatal nurse practitioner, neonatal specialty trained registered nurse, or neonatal specialty trained respiratory therapist.
- III. Transport team members should have the collective expertise sufficient to provide, if necessary:
  - A. observation of the neonate throughout the transport
  - B. monitoring of body temperature, respiratory status, and cardiovascular status
  - C. delivery and monitoring of oxygen therapy
  - D. advanced airway management and mechanical ventilation
  - E. advanced intravascular access and management skills
  - F. recognition and management of a wide variety of emergency conditions
  - G. administration of rescue and stabilization medication.
- IV. At least two members of the transport team should be current Neonatal Resuscitation Program (American Academy of Pediatrics and American Heart Association) providers. It is also recommended that team members maintain current S.T.A.B.L.E. training.
- V. Transport team members should be oriented to the transport vehicle and use of transport equipment. All transport team members should follow state and national standards.
- VI. The knowledge and skills required for a nurse to perform neonatal transport are listed in the most current edition of Educational Objectives for Nurses, Levels I, II, III Neonatal Transport Nurses, Tennessee Perinatal Care System, Tennessee Department of Health (Appendix I).

## NEONATAL SPECIALTY TRANSPORT MODALITY

- I. Selection of the mobile neonatal intensive care unit should be based on the condition of the neonate.
- II. The vehicle used exclusively for the provision of neonatal intensive care and transportation between medical facilities shall conform with the following standards for design and construction as defined by the EMS Division of the Tennessee Department of Health (Tennessee Emergency Medical Services Statutes and Rules, Rule 1200-12-1-.02[4] Special Vehicle Requirements).
  - A. Exterior surfaces, emblems, and markings shall conform to Federal Specifications-Ambulance.
  - B. Additional markings, legends, or logos may be used to identify the provider and purpose for specific vehicles, except that no letter shall exceed 14 inches in height. Legends as "neonatal intensive care or critical care transport" may be substituted for the "Ambulance" in exterior markings.
  - C. Warning lights and siren shall be furnished in accordance with Federal Specification-Ambulance except that side floodlights shall not be required.
  - D. Vehicle crashworthiness shall be assured with roll-cage construction evidenced by compliance with the Ambulance Manufacturer's Division Standards of the Truck Body and Equipment Association or comparable construction under written statement and performance bond by the manufacturer.
  - E. Doors shall provide access to the rear and curbside of the patient compartment. Where the vertical lift distance of the patient loading area exceeds 28 inches, a ramp or electrical-hydraulic lift shall be furnished to facilitate patient loading.
  - F. Environmental systems on the unit shall meet heating/air-conditioning standards as specified in Federal Specifications-Ambulance.
  - G. Vehicle electrical systems shall be provided to furnish 110 volt AC power sufficient to sustain 3,000 watts at 60 cycles. The unit shall be equipped with a back-up power system sufficient to operate patient care equipment in the event of failure of the main power systems. The 110 volt system shall incorporate a ground fault interrupter device for protection against electrical hazards.
  - H. Patient compartment shall be so designed to provide the following:
    1. a properly secured transport incubator allowing observation from at least two sides
    2. compartments for appropriate storage of materials
    3. illumination at the primary patient care area of at least 75 foot candles
    4. safety features such as sculpted, padded, or recessed cabinet corners and latches to prevent undue injury during sudden deceleration
    5. safety devices such as a grab rail or hand strap, secured according to Federal Motor Vehicle Safety Standards for safety restraints, safety belts provided at all attendant seats, and safety restraint devices for neonates
  - I. An oxygen system with sufficient capacity to deliver a minimum continuous flow of 10 liters per minute for at least four hours must be available. The installed oxygen system shall be capable of delivering specific monitored blended oxygen concentrations.
  - J. The vehicle shall provide environmental conditions for the neonate that minimize the

risk of temperature instability and excessive noise and vibration.

- III. In the event a RWA or FWA is used for neonatal transport, the following additional precautions should be observed:
  - A. use of a licensed air medical transport program with appropriately credentialed neonatal care providers
  - B. secure fastening of transport equipment
  - C. an independent power source to allow uninterrupted and fail-safe operation of the incubator and other supporting equipment
  - D. environmental conditions for the neonate that minimize the risk of temperature instability, noise and vibration
  - E. protective headgear for crew use during transport
  - F. system for easy communication among the members of the transport team and medical control at receiving center
  - G. seat belts
  - H. seating arrangement that permits close observation and handling of the patient
- IV. Latex free equipment and supplies must be provided.

## NEONATAL TRANSPORT EQUIPMENT

- I. Organization and maintenance of neonatal transport equipment is the responsibility of the transporting facility. (See Appendix VIII)
- II. Equipment to maintain a neutral thermal environment for the neonate should include:
  - A. transport incubator
  - B. thermometer
  - C. blanket, insulating blanket or chemically activated heat pack (appropriate for neonatal use)
- III. The transport incubator should meet the following requirements:
  - A. approved by the manufacturer for use during transport and installed in the transport vehicle with crashworthy restraints.
  - B. if used in RWA and FWA it shall meet FAA requirements for crashworthiness and flammability of materials. The transport incubator and monitoring equipment should be tested by an FAA certified mechanic to assure equipment does not interfere with navigational instruments.
  - C. a heat source that requires minimal time for preheating and should maintain ambient temperature within the desired range of 29° to 36° C. The control for temperature setting should be readily accessible and easy to operate, and there should be provision for easy determination of ambient temperature. It is essential to have a fail-safe alarm system that will recognize overheating or underheating.
  - D. provide an environment in which the oxygen supply is constant and controllable.
  - E. provide unrestricted visibility of the neonate with a functional independent light source for general illumination provided in or on the incubator.
  - F. easy accessibility to the neonate resulting in minimal interference with thermal protection and oxygen supply.
  - G. safety restraint devices to secure the neonate inside the incubator that conform to Federal Motor Vehicle Safety Standards and FAA Standards.
- IV. Equipment for oxygen delivery and monitoring should include:
  - A. oxygen tanks
  - B. air tanks
  - C. pressure gauges
  - D. flowmeters (needle control valve preferred)
  - E. oxygen analyzer
  - F. oxygen blender
  - G. oxygen tubing and adapters
  - H. oxygen hood or nasal cannula
  - I. neonatal oxygen mask
  - J. neonatal resuscitation manual bag and mask with manometer or T-piece resuscitator
  - K. continuous positive airway pressure apparatus
  - L. mechanical ventilator
- V. The following guidelines are related to the use of oxygen during transport:

- A. a portable supply of oxygen and compressed air in cylinders adequate to last the entire journey with surplus to cover unexpected needs and delays should be carried. Proper restraint of these cylinders is mandatory throughout the transport. An oxygen cylinder usage chart may be used in determining the number and type of oxygen cylinders necessary during transport (Appendix III).
  - B. oxygen cylinders in use should be provided with pressure gauges and flow meters.
  - C. the ambient oxygen concentration must be monitored continuously by an oxygen analyzer.
- VI.** Devices to maintain the patency of the airway and gastric decompression must be readily available and should include:
- A. bulb syringe
  - B. regulated suction with gauge
  - C. suction catheters (#6, 8, 10 Fr)
  - D. feeding tube (#8 Fr) with a 20 mL syringe
  - F. Replogle tube (#8 Fr)
- VII.** Equipment for vital sign monitoring should include:
- A. a continuous heart rate monitor
  - B. neonatal stethoscope
  - C. body temperature monitor
  - D. noninvasive and invasive blood pressure monitoring devices
  - E. pulse oximeter for noninvasive monitoring of oxygen saturation
- VIII.** Equipment for monitoring blood glucose must be available.
- IX.** If intravenous therapy is required, an infusion pump that is portable, battery-powered, fail-safe, and calibrated to ensure accurate delivery of calculated fluid volumes must be used. Critical neonates may require multiple intravenous lines requiring additional infusion pumps.
- X.** The equipment and supplies required for resuscitation of a neonate must be available, portable and should include:
- A. Endotracheal intubation
    - 1. laryngoscope handle with blades (#00, 0, 1)
    - 2. laryngoscope spare bulbs
    - 3. laryngoscope spare batteries
    - 4. endotracheal tubes (#2.5, 3.0, 3.5, 4.0 mmID)
    - 5. neonatal resuscitation manual bag and mask or T-piece resuscitator
    - 6. pressure manometer (to monitor PIP and PEEP if manual ventilation is necessary)
    - 7. disposable stylet (#6 Fr)
    - 8. adhesive tape or commercial endotracheal tube holders
    - 9. scissors
    - 10. end tidal carbon dioxide (CO<sub>2</sub>) detector
  - B. Intravenous infusion

1. intravenous needles and catheters (#22, 23, 24, 25, 26 gauge)
  2. syringes (1, 3, 6, 12, 20, 35 mL)
  3. intravenous armboard
  4. intravenous tubing and T connector
  5. infusion device
  6. tape or site dressing
  7. site preps
  8. 3-way stopcock
  9. umbilical catheterization equipment
- C. Medications should include the following drugs or an approved therapeutic equivalent based on local program protocols.
1. adenosine
  2. ampicillin
  3. atropine
  4. calcium gluconate
  5. dextrose solution (D5W)
  6. dextrose solution (D10W)
  7. dextrose solution (D25W)
  8. dobutamine
  9. dopamine
  10. epinephrine (1:10,000)
  11. fentanyl
  12. furosemide
  13. gentamicin
  14. glucagon
  15. heparin
  16. insulin (regular)
  17. morphine sulfate
  18. naloxone hydrochloride
  19. normal saline
  20. phenobarbital
  21. prostaglandin E<sub>1</sub> (requires refrigeration)
  22. sodium bicarbonate (4.2%)
  23. sterile water
  24. surfactant (requires refrigeration)
  25. THAM or other buffer

**XI. Equipment for diagnosis and management of air leak syndrome**

- A. transilluminator
- B. chest tubes (8, 10, 12 Fr)
- C. chest tube kit
- D. 18 to 20 gauge over the needle catheters
- E. drain with one-way check valve

- F. 3-way stopcock
- G. luer lock and slip tip syringe (35/60cc)

**XII.** Equipment for handwashing and personal protection

- A. antiseptic solution or towelettes
- B. gloves
- C. full face protection or goggles and masks
- D. fluid-retardant and fluid-resistant gowns
- E. infectious waste disposal bags
- F. soiled linen disposal bags
- G. sharps box

**XIII.** Latex free equipment and supplies must be provided.

## **NEONATAL REFERRAL DOCUMENTATION**

- I. Records are essential for continuing care of the patient and evaluation of the referral process. Both referring and receiving center personnel have responsibilities to provide adequate documentation of clinical data.
  
- II. **REFERRING CENTER RESPONSIBILITIES**
  - A. The following documents should accompany the transported neonate:
    - 1. copy of complete maternal prenatal record
    - 2. copy of current maternal medical record
    - 3. copy of current neonatal medical record
    - 4. copies of any images in a format that can be reviewed by the transport team at the bedside
    - 5. copy of record of care during transport
  
  - B. The referring center should maintain a record regarding disposition of transported neonates.
  
- III. **RECEIVING CENTER RESPONSIBILITIES**
  - A. maintain a record of consultation/referral calls
  - B. maintain a record regarding the disposition of the transported neonate
  - C. send a summary of care to the referring perinatal health care providers

## EVALUATION OF NEONATAL REFERRAL PROCESS

- I. Interhospital care of the high-risk neonate requires the cooperation and coordination of many skilled persons. Outreach education efforts should include discussions of the regional referral process and can be used to reinforce cooperation and coordination.
- II. Outreach education related to transport should focus on the following objectives:
  - A. informing perinatal care and EMS providers in the region of specialized resources available to them through the perinatal network.
  - B. assisting neonatal care providers in developing their abilities to identify high-risk neonatal patients, anticipate complications, and stabilize those patients before transport.
  - C. continuing quality improvement through ongoing education of neonatal care providers and EMS providers (See Appendices I and II).
- III. Planning of the neonatal referral process requires participation of those who will use the service and those who will provide it. Criteria considered in planning and evaluating the referral process are:
  - A. availability
  - B. accessibility
  - C. responsiveness
  - D. effectiveness
  - E. safety
- IV. Referring facilities should periodically review their neonatal referrals.

# **RETURN TRANSPORT**

## **RETURN TRANSPORT**

I. Return transports occur to take patients back to their original or local hospital for further care when the problems that required initial transport have been resolved. Return transports should be recognized as an important benefit to the individual patient, family, primary care provider, and regionalized perinatal care system. Early planning of return transports is desirable.

### **II. MATERNAL RETURN TRANSPORT**

- A. The hospital to which the maternal patient is returned and the timing of the return transport are largely determined by individual patient care needs and receiving institutional capabilities.
- B. The composition of the transport team should be based on the condition of the patient.
- C. A ground ambulance is the most appropriate vehicle for the majority of maternal return transports. If an alternative form of transportation is being considered, the referring health care provider should discuss this alternative mode with the receiving physician.
- D. A telephone consultation with the receiving health care provider is necessary to initiate the return transport process and to prepare the receiving center. This consultation may aid the physician in developing a treatment plan.
- E. The physician directing the return transport is responsible for the patient during transport.
- F. Consent forms to authorize transfer, treatment, and admission to the receiving hospital must be obtained (COBRA, EMTALA; see Appendix VI).
- G. A summary of care must accompany the patient.

### **III. NEONATAL RETURN TRANSPORT**

- A. The hospital to which the neonate is returned and the timing of the return transport are largely determined by individual patient care needs and receiving institutional capabilities.
- B. A telephone consultation with the receiving health care provider is necessary to initiate the return transport process and to prepare the receiving hospital. This consultation may aid the health care provider and nursing staff in developing a treatment plan. In the event the neonate is returned to a hospital other than the original referring hospital, the original referring health care provider must be notified.
- C. The mode of transport, composition of the transport team, and equipment needs should be based on the condition of the neonate and other factors such as distance and weather conditions.
- D. The physician directing the return transport is responsible for the patient during transport.
- E. Consent forms to authorize transfer, treatment, and admission to the receiving center must be obtained.
- F. The parents should be encouraged to visit and become familiar with the receiving center nursery prior to the return transport.
- G. The transport team should communicate with receiving center personnel regarding the estimated time of arrival.
- H. On admission of the neonate to the receiving center, the transport team should

communicate with receiving center personnel regarding the neonate's history, events during transport, and current status (Appendix IV).

- I. A summary of care should accompany the patient.
- J. Periodic communication between referring and receiving hospitals should be maintained (Appendix V).

## **APPENDICES**

## APPENDIX I

### EDUCATIONAL OBJECTIVES FOR NEONATAL TRANSPORT NURSES

Educational Objectives for Nurses, Levels I, II, III, Neonatal Transport Nurses (the most recent edition), Tennessee Perinatal Care System, Tennessee Department of Health.

The nurse caring for neonatal patients during transport should be able to meet the objectives listed for each of the following categories:

#### I. PROBLEMS OF PREGNANCY, FETAL DEVELOPMENT, LABOR AND DELIVERY

- A. Obtain from a referring caretaker reports of all tests done to determine fetal gestational age and well-being.
- B. Utilize data from the maternal/neonatal history as a basis for anticipating problems, planning, and implementing care during transport.
- C. Provide for a receiving caretaker, maternal and neonatal data which give adequate history of problems resulting from pregnancy, labor, and delivery, as well as treatment provided.

#### II. RESUSCITATION OF THE NEONATE

- A. Provide for a receiving caretaker an accurate record of required resuscitative procedures and the neonate's physiological responses.

#### III. PHYSICAL ASSESSMENT OF THE NEWBORN

- A. Collaborate with other transport team members in obtaining a thorough physical assessment prior to transport.
- B. Describe and initiate an assessment plan during transport that will identify infant problems when they are most amenable to intervention.
- C. Provide for a receiving physician a complete record of physical assessment, which includes information from the referring care providers, as well as the transport staff.

#### IV. THERMOREGULATION

- A. Explain the effect of environmental factors, e.g., humidity, ambient temperature, altitude, and velocity of air flow on the thermal status of the neonate.
- B. Describe safe methods of maintaining, increasing, and decreasing a neonate's temperature in a transport situation.
- C. Provide a receiving caretaker with a thorough history of the infant's thermoregulation problems, treatment of these problems, and infant responses to intervention prior to and during transport.

#### V. NUTRITIONAL REQUIREMENTS OF THE NEONATE

- A. Describe the effects of speed, acceleration, and deceleration on gastrointestinal motility and sphincter control.
- B. Describe safe means of providing infant nutrition in a variety of transport situations.

- C. Obtain from a referring caretaker an accurate nutritional record for the receiving caretaker.

## **VI. INTRAVASCULAR THERAPY**

- A. Describe and utilize safe, efficient measures to limit the effects of transport on intravascular therapy.
- B. Prepare fluids and blood products that may be required during transport.
- C. Record for a receiving caretaker an accurate summary of fluid and blood products infused prior to and during transport.

## **VII. MEDICATION ADMINISTRATION**

- A. Prepare medications that may be required during transport.
- B. Provide for a receiving caretaker an accurate record of medications used prior to and during transport and the neonate's responses.

## **VIII. FLUID, ELECTROLYTE, AND ACID-BASE BALANCE**

- A. Describe the effects of marked changes in humidity, velocity, and pressure on insensible fluid loss, and measures to limit these effects.
- B. Obtain and record an accurate summary of fluid, electrolyte and acid-base status prior to and during transport.

## **IX. RESPIRATORY DISORDERS OF THE NEWBORN**

- A. Describe the effects of altering atmospheric pressure, altitude, temperature, and humidity on neonatal respiratory function, and discuss nursing measures to minimize these effects.
- B. Select and utilize respiratory measures, pharmacologic agents, intravenous orders, and infant positioning to assist in lessening or preventing the disorders listed above.
- C. Obtain an accurate history of respiratory status and respiratory support provided prior to transport and develop an ongoing record of assessment, evaluation, and respiratory support for the receiving center.

## **X. RESPIRATORY SUPPORT SYSTEMS**

- A. Determine the settings to be used when the infant is switched from:
  - 1. A pressure-cycled to a time-cycled/pressure-limited ventilator;
  - 2. A time-cycled/pressure-limited ventilator to a pressure-cycled ventilator;
  - 3. A volume-cycled ventilator to a pressure-cycled ventilator.
- B. Set up and correctly utilize respiratory support and monitoring equipment used during transport.

## **XI. HEMATOLOGIC DISORDERS OF THE NEWBORN**

- A. Obtain appropriate consent for the administration of blood products.

- B. Collaborate with the Transport Team Leader in obtaining reports or specimens for a hematologic data base, including information on the treatment of these disorders prior to and during transport.
- C. Collaborate with the referral center and/or transport service in obtaining blood or blood products which may be required during transport to the receiving center.
- D. Provide for a receiving caretaker an accurate hematologic history, including treatment during transport.

## **XII. GASTROINTESTINAL PROBLEMS OF THE NEWBORN**

- A. Identify the special techniques and measures required to limit the side effects of gastrointestinal obstructions and/or abdominal wall defects during transport.
- B. Provide for a receiving caretaker, a history of gastrointestinal function, treatment, and neonatal response prior to and during transport.

## **XIII. PERINATAL INFECTION**

- A. Collaborate with the Transport Team Leader in providing the different components of a septic work-up, in a safe and timely manner.
- B. Develop and implement procedures which will enhance prevention of infection in transport situations.
- C. Obtain from a referring caretaker and provide for the receiving caretaker a history which identifies a neonate's risk of infection and antibiotic therapy provided.

## **XIV. CARDIAC DISORDERS OF THE NEONATE**

- A. Design and implement a plan of care that will provide maximum protection from hypoxic damage for the infant who has cardiac disorders.
- B. Provide for a receiving caretaker a thorough report of cardiovascular problems, assessment, treatment, and neonatal condition prior to and during transport.

## **XV. PARENT-INFANT RELATIONSHIPS**

- A. Describe and utilize measures which will enhance a positive relationship between parents and health care personnel in the referring and receiving centers.
- B. Describe the potential effects of transport on the development of a positive parent-infant relationship.
- C. Describe and utilize measures that will minimize the negative effects of transport on parent-infant bonding.
- D. Provide for a receiving caretaker a report of significant parent, neonate, and staff interactions.

## **XVI. REFERRING-RECEIVING CARETAKER RELATIONSHIPS**

- A. When given a report by a referring caretaker, anticipate and rapidly request information necessary to provide continuous expert care.
- B. Collaborate with other nurses in the perinatal region in developing transport plans that provide comprehensive, continuous, and expert care.

- C. Describe the general types of services available in Level I, II, and III newborn facilities.
- D. Identify and communicate effectively the attributes and limitations of Level I, II, and III facilities in the region.
- E. Describe, utilize, and communicate to others appropriate procedures for initiating consultation, referral, and transport.
- F. Describe and prepare the written records required prior to transport.
- G. Identify and evaluate communication patterns in the transport region.
- H. Seek and accept constructive evaluation of the referral process from nurses in Level I, II, and III facilities.
- I. Utilize constructive criticism and effective communication skills as a basis for improving individual care, improving continuity of expert care within the regional center, and improving care of infants referred to and received from other facilities.

## **XVII. TRANSPORT SAFETY**

- A. Describe those factors that must be considered in the selection of a vehicle and professional personnel for transport.
- B. Describe and utilize effective techniques for securing transport equipment and compressed medical gas tanks in transport vehicles.
- C. Determine adequacy of illumination in transport vehicles.
- D. Provide continuous visibility of the infant, support equipment, and monitors during transport.
- E. Determine that space available in the transport vehicle is adequate for safe emergency intervention during transport.
- F. Describe briefly the effects of vibration and sound level on the infant in transit and develop a plan (ear muffs) to diminish these effects.
- G. Determine the adequacy of power sources to assure uninterrupted power availability during transport.
- H. State the potential hazards of vehicle acceleration, deceleration, and speed on the transported infant, and take appropriate measures to limit their occurrence, including an appropriate restraint system.
- I. Determine and provide an adequate supply of oxygen and compressed air required for transport.
- J. Describe and utilize effective methods for testing equipment function prior to transport.
- K. Identify and provide the life support and monitoring equipment and supplies necessary for transport.
- L. Implement a plan which provides for replacement, cleaning, and maintenance of transport vehicle, equipment, and supplies.
- M. Describe to others and utilize appropriate steps for stabilizing the infant prior to transport.
- N. Utilize the vehicle communication system effectively in obtaining consultation from

other professional personnel during transport.

- O. Communicate an infant assessment which will assure adequate professional support and equipment upon the arrival of the transported infant at the receiving center.
- P. Maintain records, which can be readily utilized to evaluate the effectiveness of the transport system.
- Q. Assist in evaluation and implement measures to improve the transport process.

## APPENDIX II

### Education Requirements for Emergency Medical Services Personnel

National Standard Curricula for Emergency Medical Services providers are available on the web site of the National Highway Traffic Safety Administration, United States Department of Transportation ([www.nhtsa.gov](http://www.nhtsa.gov)). Training curricula are available for First Responders, EMTs, and Paramedics. On the web site under Traffic Safety, query for National Standard Curricula.

See *Emergency Medical Technician: Basic Refresher Curriculum*, Module VI: Obstetrics, Infants, and Children, and Appendix A: USDOT Curriculum Objectives for Emergency Medical Technician: Basic Refresher, which lists objectives from the entire EMT–Basic National Standard Curriculum (<http://www.nhtsa.dot.gov/people/injury/ems/pub/basicref.pdf>).

#### The training objectives for Paramedics related to Neonatology are:

United States Department of Transportation National Highway Traffic Safety Administration  
**Paramedic: National Standard Curriculum**  
**NEONATOLOGY**

#### **TERMINAL OBJECTIVE**

At the completion of this unit, the paramedic student will be able to integrate pathophysiological principles and assessment findings to formulate a field impression and implement a treatment plan for a neonatal patient.

#### **COGNITIVE OBJECTIVES**

At the completion of this unit, the paramedic student will be able to:

- 6-1.2 Define the term newborn. (C-1)
- 6-1.3 Define the term neonate. (C-1)
- 6-1.4 Identify important antepartum factors that can affect childbirth. (C-1)
- 6-1.5 Identify important intrapartum factors that can term the newborn high risk. (C-1)
- 6-1.6 Identify the factors that lead to premature birth and low birth weight newborns. (C-1)
- 6-1.7 Distinguish between primary and secondary apnea. (C-3)
- 6-1.8 Discuss pulmonary perfusion and asphyxia. (C-1)
- 6-1.9 Identify the primary signs utilized for evaluating a newborn during resuscitation. (C-1)
- 6-1.10 Formulate an appropriate treatment plan for providing initial care to a newborn. (C-3)
- 6-1.11 Identify the appropriate use of the APGAR score in caring for a newborn. (C-1)
- 6-1.12 Calculate the APGAR score given various newborn situations. (C-3)
- 6-1.13 Determine when ventilatory assistance is appropriate for a newborn. (C-1)
- 6-1.14 Prepare appropriate ventilation equipment, adjuncts, and technique for a newborn. (C-1)
- 6-1.15 Determine when chest compressions are appropriate for a newborn. (C-1)
- 6-1.16 Discuss appropriate chest compression techniques for a newborn. (C-1)
- 6-1.17 Assess patient improvement due to chest compressions and ventilations. (C-1)
- 6-1.18 Determine when endotracheal intubation is appropriate for a newborn. (C-1)

- 6-1.19 Discuss appropriate endotracheal intubation techniques for a newborn. (C-1)
- 6-1.20 Assess patient improvement due to endotracheal intubation. (C-1)
- 6-1.21 Identify complications related to endotracheal intubation for a newborn. (C-1)
- 6-1.22 Determine when vascular access is indicated for a newborn. (C-1)
- 6-1.23 Discuss the routes of medication administration for a newborn. (C-1)
- 6-1.24 Determine when blow-by oxygen delivery is appropriate for a newborn. (C-1)
- 6-1.25 Discuss appropriate blow-by oxygen delivery devices and technique for a newborn. (C-1)
- 6-1.26 Assess patient improvement due to assisted ventilations. (C-1)
- 6-1.27 Determine when an orogastric tube should be inserted during positive-pressure ventilation. (C-1)
- 6-1.28 Discuss the signs of hypovolemia in a newborn. (C-1)
- 6-1.29 Discuss the initial steps in resuscitation of a newborn. (C-1)
- 6-1.30 Assess patient improvement due to blow-by oxygen delivery. (C-1)
- 6-1.31 Discuss the effects maternal narcotic usage has on the newborn. (C-1)
- 6-1.32 Determine the appropriate treatment for the newborn with narcotic depression. (C-1)
- 6-1.33 Discuss appropriate transport guidelines for a newborn. (C-1)
- 6-1.34 Determine appropriate receiving facilities for low and high risk newborns. (C-1)
- 6-1.35 Describe the epidemiology, including the incidence, morbidity/ mortality, risk factors and prevention strategies for meconium aspiration. (C-1)
- 6-1.36 Discuss the pathophysiology of meconium aspiration. (C-1)
- 6-1.37 Discuss the assessment findings associated with meconium aspiration. (C-1)
- 6-1.38 Discuss the management/ treatment plan for meconium aspiration. (C-1)
- 6-1.39 Describe the epidemiology, including the incidence, morbidity/ mortality, risk factors and prevention strategies for apnea in the neonate. (C-1)
- 6-1.40 Discuss the pathophysiology of apnea in the neonate. (C-1)
- 6-1.41 Discuss the assessment findings associated with apnea in the neonate. (C-1)
- 6-1.42 Discuss the management/ treatment plan for apnea in the neonate. (C-1)
- 6-1.43 Describe the epidemiology, pathophysiology, assessment findings, management/ treatment plan for diaphragmatic hernia. (C-1)
- 6-1.44 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for bradycardia in the neonate. (C-1)
- 6-1.45 Discuss the pathophysiology of bradycardia in the neonate. (C-1)
- 6-1.46 Discuss the assessment findings associated with bradycardia in the neonate. (C-1)
- 6-1.47 Discuss the management/ treatment plan for bradycardia in the neonate. (C-1)
- 6-1.48 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for premature infants. (C-1)
- 6-1.49 Discuss the pathophysiology of premature infants. (C-1)
- 6-1.50 Discuss the assessment findings associated with premature infants. (C-1)
- 6-1.51 Discuss the management/ treatment plan for premature infants. (C-1)
- 6-1.52 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for respiratory distress/ cyanosis in the neonate. (C-1)
- 6-1.53 Discuss the pathophysiology of respiratory distress/ cyanosis in the neonate. (C-1)
- 6-1.54 Discuss the assessment findings associated with respiratory distress/ cyanosis in the neonate. (C-1)

- 6-1.55 Discuss the management/ treatment plan for respiratory distress/ cyanosis in the neonate. (C-1)
- 6-1.56 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for seizures in the neonate. (C-1)
- 6-1.57 Discuss the pathophysiology of seizures in the neonate. (C-1)
- 6-1.58 Discuss the assessment findings associated with seizures in the neonate. (C-1)
- 6-1.59 Discuss the management/ treatment plan for seizures in the neonate. (C-1)
- 6-1.60 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for fever in the neonate. (C-1)
- 6-1.61 Discuss the pathophysiology of fever in the neonate. (C-1)
- 6-1.62 Discuss the assessment findings associated with fever in the neonate. (C-1)
- 6-1.63 Discuss the management/ treatment plan for fever in the neonate. (C-1)
- 6-1.64 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for hypothermia in the neonate. (C-1)
- 6-1.65 Discuss the pathophysiology of hypothermia in the neonate. (C-1)
- 6-1.66 Discuss the assessment findings associated with hypothermia in the neonate. M (C-1)
- 6-1.67 Discuss the management/ treatment plan for hypothermia in the neonate. (C-1)
- 6-1.68 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for hypoglycemia in the neonate. (C-1)
- 6-1.69 Discuss the pathophysiology of hypoglycemia in the neonate. (C-1)
- 6-1.70 Discuss the assessment findings associated with hypoglycemia in the neonate. (C-1)
- 6-1.71 Discuss the management/ treatment plan for hypoglycemia in the neonate. (C-1)
- 6-1.72 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for vomiting in the neonate (C-1)
- 6-1.73 Discuss the pathophysiology of vomiting in the neonate. (C-1)
- 6-1.74 Discuss the assessment findings associated with vomiting in the neonate. (C-1)
- 6-1.75 Discuss the management/ treatment plan for vomiting in the neonate. (C-1)
- 6-1.76 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for diarrhea in the neonate. (C-1)
- 6-1.77 Discuss the pathophysiology of in diarrhea the neonate. (C-1)
- 6-1.78 Discuss the assessment findings associated with diarrhea in the neonate. (C-1)
- 6-1.80 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for common birth injuries in the neonate. (C-1)
- 6-1.81 Discuss the pathophysiology of common birth injuries in the neonate. (C-1)
- 6-1.82 Discuss the assessment findings associated with common birth injuries in the neonate. (C-1)
- 6-1.83 Discuss the management/ treatment plan for common birth injuries in the neonate. (C-1)
- 6-1.84 Describe the epidemiology, including the incidence, morbidity/ mortality and risk factors for cardiac arrest in the neonate. (C-1)
- 6-1.85 Discuss the pathophysiology of cardiac arrest in the neonate. (C-1)
- 6-1.86 Discuss the assessment findings associated with cardiac arrest in the neonate. (C-1)
- 6-1.87 Discuss the management/ treatment plan for cardiac arrest in the neonate. (C-1)
- 6-1.88 Discuss the pathophysiology of post arrest management of the neonate. (C-1)
- 6-1.89 Discuss the assessment findings associated with post arrest situations in the neonate. (C-1)

6-1.90 Discuss the management/ treatment plan to stabilize the post arrest neonate. (C-1)

### **AFFECTIVE OBJECTIVES**

At the completion of this unit, the paramedic student will be able to:

- 6-1.91 Demonstrate and advocate appropriate interaction with a newborn/ neonate that conveys respect for their position in life. (A-3)
- 6-1.92 Recognize the emotional impact of newborn/ neonate injuries/ illnesses on parents/ guardians. (A-1)
- 6-1.93 Recognize and appreciate the physical and emotional difficulties associated with separation of the parent/ guardian and a newborn/ neonate. (A-3)
- 6-1.94 Listen to the concerns expressed by parents/ guardians. (A-1)
- 6-1.95 Attend to the need for reassurance, empathy, and compassion for the parent/ guardian. (A-1)

### **PSYCHOMOTOR OBJECTIVES**

At the completion of this unit, the paramedic student will be able to:

- 6-1.96 Demonstrate preparation of a newborn resuscitation area. (P-2)
- 6-1.97 Demonstrate appropriate assessment technique for examining a newborn. (P-2)
- 6-1.98 Demonstrate appropriate assisted ventilations for a newborn. (P-2)
- 6-1.99 Demonstrate appropriate endotracheal intubation technique for a newborn. (P-2)
- 6-1.100 Demonstrate appropriate meconium aspiration suctioning technique for a newborn. (P-2)
- 6-1.101 Demonstrate appropriate insertion of an orogastric tube. (P-2)
- 6-1.102 Demonstrate needle chest decompression for a newborn or neonate. (P-2)
- 6-1.103 Demonstrate appropriate chest compression and ventilation technique for a newborn. (P-2)
- 6-1.104 Demonstrate appropriate techniques to improve or eliminate endotracheal intubation complications. (P-2)
- 6-1.105 Demonstrate vascular access cannulation techniques for a newborn. (P-2)
- 6-1.106 Demonstrate the initial steps in resuscitation of a newborn. (P-2)
- 6-1.107 Demonstrate blow-by oxygen delivery for a newborn. (P-2)

## APPENDIX III

### OXYGEN CYLINDER USAGE DURING TRANSPORT

#### OXYGEN CYLINDERS: DURATION OF FLOW

##### **SIMPLE FORMULA:**

Gauge pressure in psi (pounds per square inch) minus the safe residual pressure (always 200 psi) times the constant (see list below) divided by the flow rate in liters per minute = duration of flow in minutes.

##### **CYLINDER CONSTANTS**

D = 0.16    G = 2.41  
E = 0.28    H = 3.14  
M = 1.56    K = 3.14

##### **EXAMPLE**

Determine the life of an M cylinder that has a pressure of 2000 psi displayed on the pressure gauge and a flow rate of 10 liters per minute.

$$\frac{(2000-200) \times 1.56}{10} = \frac{2808}{10} = 280.8 \text{ minutes}$$

#### OXYGEN SUPPLY AND REGULATORS

Oxygen is supplied either as a compressed gas or as a liquid. Compressed gaseous oxygen is stored in an aluminum or steel tank in 400 liter (D), 660 liter (E), or 3,450 liter (M) volumes. To calculate how long the oxygen will last:

$$\text{Tank life in minutes} = (\text{tank pressure in psi} \times 0.28) \div \text{liters per minute}$$

## Appendix IV

### An Example of a Neonatal Return Transport Worksheet

#### NURSING TRANSFER SHEET BACK TRANSPORT

Date \_\_\_\_\_  
 RN \_\_\_\_\_ NNP \_\_\_\_\_  
 Driver \_\_\_\_\_ Mode of transport \_\_\_\_\_  
 Receiving Hosp/phone # \_\_\_\_\_  
 Receiving Nurse \_\_\_\_\_  
 Receiving MD/phone # \_\_\_\_\_

Addressograph / Label

Patient Information

DOB \_\_\_\_\_  
 Birth Place \_\_\_\_\_  
 BW / GA \_\_\_\_\_  
 Weight / race \_\_\_\_\_  
 Diagnosis \_\_\_\_\_  
 Parents \_\_\_\_\_  
 \_\_\_\_\_  
 Address \_\_\_\_\_  
 \_\_\_\_\_  
 Phone # \_\_\_\_\_  
 \_\_\_\_\_

Nursing Information

Bed temp \_\_\_\_\_  
 Type Feeding \_\_\_\_\_  
 Route \_\_\_\_\_  
 Amount \_\_\_\_\_  
 Last feed \_\_\_\_\_  
 Feeding problems: \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 Recipe: \_\_\_\_\_  
 \_\_\_\_\_

History / Needs

Resp \_\_\_\_\_  
 Apnea \_\_\_\_\_  
 \_\_\_\_\_  
 Cardiac \_\_\_\_\_  
 Hematology \_\_\_\_\_  
 \_\_\_\_\_  
 Immunizations \_\_\_\_\_  
 HUS \_\_\_\_\_  
 Audiology \_\_\_\_\_  
 Other needs \_\_\_\_\_

Check:

\_\_\_\_\_ Transport consent  
 \_\_\_\_\_ Discharge summary  
 \_\_\_\_\_ Transfer orders  
 \_\_\_\_\_ Newborn screen/PKU  
 \_\_\_\_\_ Personal items/milk  
 \_\_\_\_\_  
 \_\_\_\_\_ Depart Referring Hosp  
 \_\_\_\_\_ Arrive Receiving Hosp  
 \_\_\_\_\_ Depart Receiving Hosp  
 \_\_\_\_\_ Arrive Vanderbilt

IVF # 1:

Type \_\_\_\_\_  
 Route \_\_\_\_\_  
 Rate \_\_\_\_\_  
 # 2: \_\_\_\_\_  
 \_\_\_\_\_

FU appt

\_\_\_\_\_

	Medication	Dosage	Route	Frequency	Last dose
1.	_____				
2.	_____				
3.	_____				
4.	_____				

Transport Record								Assessment / Observations	
Time								ID band # / site:	
Ax / Bed T									
HR									
RR									
BP / Mean									
IMV									
PIP / PEEP									
FiO <sub>2</sub> / sats									
IVF / amt infused									

Signature of Transport Nurse: \_\_\_\_\_

## Appendix V

### An Example of a Neonatal Return Transport Follow-up Form

**PLEASE RETURN TO** (REFERRING FACILITY)

Addressograph / Label

**Follow-up Information on  
Back-Transported Infants**

Infant's name: \_\_\_\_\_  
 Date of birth: \_\_\_\_\_ Birth wt. \_\_\_\_\_ G.A. \_\_\_\_\_  
 Back transport date: \_\_\_\_\_ Discharge diagnosis: \_\_\_\_\_  
 Receiving hospital: \_\_\_\_\_ Receiving physician: \_\_\_\_\_  
 Current wt. \_\_\_\_\_ Length \_\_\_\_\_ H.C. \_\_\_\_\_ P.M.A. \_\_\_\_\_ O<sub>2</sub> requirement \_\_\_\_\_

**Please fill in upon discharge home and return to address on back or fax.**

Date of discharge: \_\_\_\_\_ Discharge wt. \_\_\_\_\_  
 PCP or health clinic: \_\_\_\_\_  
 (Current parents/custodians) Name: \_\_\_\_\_  
 Street address: \_\_\_\_\_  
 City/State/Zip: \_\_\_\_\_  
 Area code/ Phone #: \_\_\_\_\_

**Clinical course in your institution prior to discharge:**

Total days on O<sub>2</sub> after transport? \_\_\_\_\_ BPD appt. (insert phone #) if home on O<sub>2</sub>? Y/N

	Yes	No	Follow-up appt (y/n)
Ophthalmology exams?			
Head ultrasounds?			
Audiology tests?			
Synagis injection?			

**Please check all that apply at discharge:**

<b>Equipment:</b> Oxygen <input type="checkbox"/> Oximeter <input type="checkbox"/> Monitor <input type="checkbox"/> Suction <input type="checkbox"/> Other: <input type="checkbox"/>	<b>Medications:</b> Vitamins <input type="checkbox"/> Diuretics <input type="checkbox"/> Antireflux <input type="checkbox"/> Antibiotics <input type="checkbox"/> Other: <input type="checkbox"/>	<b>Feeding problems:</b> GE Reflux <input type="checkbox"/> Gastrostomy <input type="checkbox"/> Other: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<b>Referrals:</b> TEIS <input type="checkbox"/> KY First Steps <input type="checkbox"/> HUGS <input type="checkbox"/> Other: <input type="checkbox"/> <input type="checkbox"/>
--	--	---	---

**Comments:**

**FAX # (insert phone #)**  
**Attn: (insert name)**  
 (Or mail to address on reverse side)

## APPENDIX VI

### COBRA/EMTALA STATUTE: 42 USC 1395

There are a variety of resources for obtaining the COBRA/EMTALA Statute in its entirety. This information was obtained from the web site [www.medlaw.com](http://www.medlaw.com).

The COBRA/EMTALA Statute was developed in response to actions construed as patient "dumping" related to reimbursement source. The EMTALA portion relates to the transfer and medical treatment of women in active labor. The components delineate the circumstances under which an individual may be transferred to another medical care facility and the steps to be undertaken for stabilization and treatment prior to transfer.

This federal law has been in effect for a number of years. All health care facilities have been made aware of these regulations and have programs in place to address these situations. Facilities that do not abide by these regulations are subject to significant monetary sanctions.

The Transportation guidelines have been written with the understanding that all facilities will abide by the federal regulations of COBRA/EMTALA. It is the intent of the subcommittee that all facilities will abide by COBRA/EMTALA regulations.

## Appendix VII

### Neonatal Resuscitation Equipment and Supplies

#### Suction equipment:

- Bulb syringe
- Mechanical suction and tubing
- Suction catheters, 5F or 6F, 8F, 10F, 12F or 14F
- 8F feeding tube and 20-mL syringe
- Meconium aspirator

#### Bag-and-mask equipment:

- Device for delivering positive pressure ventilation, capable of delivering 90% to 100% oxygen
- Face masks, newborn and premature sizes (cushioned-rim masks preferred)
- Oxygen source with flowmeter (flow rate up to 10 L/min) and tubing

#### Intubation equipment:

- Laryngoscope with straight blades, No. 0 (preterm) and No. 1 (term)
- Extra bulbs and batteries for laryngoscope
- Endotracheal tubes, 2.5, 3.0, 3.5, 4.0 mm internal diameter (ID)
- Stylet (optional)
- Scissors
- Tape or securing device for endotracheal tube
- Alcohol sponges
- CO<sub>2</sub> detector or capnograph
- Laryngeal mask airway (optional)

#### Medications:

- Epinephrine 1:10,000 (0.1 mg/mL) 3 mL or 10 mL ampules
- Isotonic crystalloid (normal saline or Ringer's lactate) for volume expansion – 100 or 250 mL
- Sodium bicarbonate 4.2% (5 mEq/10mL) – 10-mL ampules
- Naloxone hydrochloride 0.4 mg/mL (1 mL ampules), or 1.0 mg/mL (2 mL ampules)
- Dextrose 10%, 250 mL
- Normal saline for flushes
- Umbilical vessel catheterization supplies
  - Sterile gloves
  - Scalpel or scissors
  - Antiseptic prep solution
  - Umbilical tape
  - Umbilical catheters, 3.5F, 5F
  - Three-way stopcock
- Syringes, 1, 3, 5, 10, 20, 50 mL
- Needles, 25, 21, 18 gauge, or puncture device for needleless system

#### Miscellaneous

- Gloves and appropriate personal protection
- Radiant warmer or other heat source
- Firm, padded resuscitation surface
- Clock with second hand (timer optional)
- Warmed linens
- Stethoscope (neonatal head preferred)
- Tape, ½ or ¾ inch
- Cardiac monitor and electrodes or pulse oximeter and probe (optional for delivery room)
- Oropharyngeal airways (0, 00, and 000 sizes or 30, 40, and 50mm lengths)

**For very preterm babies (optional):**

- Compressed air source
- Oxygen blender to mix oxygen and compressed air
- Pulse oximeter and oximeter probe
- Reclosable, food-grade plastic bag (1-gallon size) or plastic wrap
- Chemically activated warming pad
- Laryngoscope with straight blade, No.00
- Transport incubator to maintain baby's temperature during move to the nursery

## Appendix VIII

### TENNESSEE EMERGENCY MEDICAL SERVICES EQUIPMENT AND SUPPLIES AMBULANCE REQUIREMENTS—2006 PER EMS RULE 1200-12-1-.03, EFFECTIVE JANUARY 30, 2006

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
<b>OS OXYGEN SYSTEMS</b>				
Bag Valve Resuscitator - Adult	bag-volume of 1,600 milliliters	-	1 + spare	Disposable preferred
Bag Valve Resuscitator- Pediatric	bag-volume of 450 milliliters	-	1 + spare	Disposable preferred
Resuscitation Mask - Adult			1 + spare	Disposable preferred
Resuscitation Mask - Child			1 + spare	Disposable preferred
Resuscitation Mask - Infant			1 + spare	Disposable preferred
Oropharyngeal airways	Guedel or Berman	Adult S. M. L Child, Infant	Adult S. M. L Child, Infant	Disposable
Nasopharyngeal airways		5 sizes		Non Latex- PVC preferred
Dual lumen airway device	Combi-tube or PTL	Adult	1	Single use device
End tidal CO <sub>2</sub> Detector	Colorimetric or Numeric	Adult and pediatric	1 each	Capnometry preferred; Spare disposables recommended
Oxygen Supply tubing	Minimum 48 inch lengths		2 sets	
Adult Non-Rebreathing Oxygen Mask	For delivery of Oxygen at High Concentration	Tubing may be integrated or separate	2	
Pediatric Non-Rebreathing Oxygen Mask	For delivery of Oxygen at High Concentration		2	
Infant Medium Concentration Oxygen Mask	For delivery of Oxygen at Medium Concentration		2	
Adult Nasal Cannula	For delivery of Oxygen at Low Concentration		2	
Minimum 2,000 liter on board Oxygen Supply			1	Cylinders restrained in approved manner
Regulator and flow meters; Two distribution outlets	Pressure reducer to 50 psi piped to at least two distribution outlets	Flow meter Minimum range of 2-15 lpm	2 flow meters in patient compartment	
Minimum 300 Liter D cylinder	Portable oxygen		1 + spare	
Portable Oxygen Regulator	Pressure reducer to 50 psi, flow valve or flow meter	Minimum range of 2-15 lpm	1	Non-gravity dependent

<b>ADVANCED AIRWAY MANAGEMENT</b>				
Laryngoscope handle - Adult	With batteries		Spare batteries	
Laryngoscope handle - Pediatric	With batteries		Spare batteries	
Laryngoscope blade -infant	Straight Blade	0	1	
Laryngoscope blade - infant	Straight Blade	1	1	
Laryngoscope blade -child	Straight Blade	2	1	Note: Adds blades
Laryngoscope blade - child	Curved Blade	2	1	

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
Laryngoscope blade – adult	Straight Blade	3	1	
Laryngoscope blade- adult	Curved Blade	3	1	
Laryngoscope blade-adult	Straight Blade	4	1	
Laryngoscope blade	Curved Blade	4	1	
Endotracheal tubes-	Uncuffed	2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0	8 (1 each size) Note: Increases inventory from previous rules	Sterile for Single patient use / disposable
Endotracheal tubes	Cuffed	6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5	7 (1 each size) Note: Increases inventory from previous rules	Sterile for Single patient use/disposable
Sterile surgical lubricant	Water soluble gel	≥ 2 mL each	6 packets	Or tube with QS.
Adult stylet			1	
Pediatric stylet			1	
Magill Forceps	Adult		1	
Magill Forceps	Pediatric		1	
Esophageal Detection Device			1	Bulb type preferred

#### SUCTION DEVICES

Installed suction- with vacuum gauge, a control, and collection Per Federal Specification. 3.12.3	Minimum 1,00 mL Collection canister	≥ 300 mm/Hg vacuum	1	
Suction tubing	Six feet in length		2	
Meconium Aspirator	allowing direct connection of suction to the endotracheal tube			
Portable Suction Device w/powered pump Per Federal Specification 3.12.4	A collection bottle of at 500 milliliters shall be provided		1	disposable collection bottle preferred
Suction tubing	two feet or more in length	≥.375 I.D. preferred	2	Shorter tubing has faster draw down time
Suction catheters		6, 8, 10, 14 and 16 French ga.	2 each	Packaged; Sterile
Rigid Suction tips			2	Yankauer -type

#### DIAGNOSTIC AND ASSESSMENT DEVICES

Sphygmomanometer with inflation bulb and gauge with			1	
Adult blood pressure cuff			1	
Pediatric blood pressure cuff			1	
Adult large/ thigh blood pressure cuff			1	
Stethoscope			1	may be personal equipment, per service policy
Bandage Shears			1	may be personal equipment, per service policy
Pulse Oximeter	With adult and pediatric probes		1	Effective 3/1/2006

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
<b>BANDAGES AND DRESSINGS</b>				
Surgical adhesive tape	at least one inch in width		2 rolls	
Conforming gauze roller bandage	at least three inches in width	Comparable to Kerlix or Kling®	6 rolls	Clean or sterile supply allowed
Triangular Bandages	For slings, cravat bandages, or splint ties	base ≥ 42 inches	6	
Sterile gauze dressings 4" x 4"	Gauze dressing or sponges	4" by 4"	25 individual pads	Acceptable in packs of 2
Composite pad sterile compresses	Comparable to ABD pads	Typical sizes: 4" x 8" 8" x 10"; or 5" by 9"	8	
Sterile occlusive dressings	white petrolatum coated gauze or plastic membrane film	at least 3" by 3"	2	Aluminum foil or plastic bags not acceptable as substitute
Burn sheets	separately packaged, sterile or clean	at least 60 by 60 inches	2	Sterile cotton (cloth) sheets preferred
Irrigation fluids	Sterile Saline solution or sterile water	sufficient to supply 2000 milliliters		Plastic Containers required

<b>IMMOBILIZATION DEVICES</b>				
Long spinal immobilization devices or backboards	whole body splints, or approved devices capable of immobilizing a patient with suspected spinal injuries		2	
Straps or restraints	To immobilize a patient at or about the chest, pelvis, and knees		6 straps or 2 ≥ six point Spider-device	
Short spinal immobilization device	shall provide spinal immobilization for the seated patient		One device	with affixed padding and straps
Case or carrier bag for above	To maintain in clean condition		1	
C-Spine or Head Immobilizers	To prevent lateral head movement of the restrained patient.		2 sets or 4 blocks	With straps or restraint materials
Cervical Immobilization Collars	To prevent head and neck movement of the restrained patient.	Pediatric/Infant Small Adult Medium Adult Large Adult	at least two adult collars in each size range and two pediatric/infant collars	Combinations of adjustable-type collars are acceptable to provide at least two adult and two pediatric/infant collars
Upper extremity splints	Fabricated splints for immobilization of arm injuries	≥ 15" in length	2 splints or two sets of board splints	Capability to immobilize simultaneously two arms
Lower extremity splints	Fabricated splints for immobilization of leg injuries	Fabricated splints in assorted sizes; for board splints ≥ 36" in length	at least two devices or two sets of board splints	Capability to immobilize simultaneously two legs

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
Lower extremity traction splints	Fabricated splints for immobilization of femoral fractures	Commercial devices	At least two devices or one device designed to immobilize both legs	Capability to immobilize simultaneously two legs

PATIENT CARE SUPPLIES				
Emesis basin or suitable substitute	Containers for human waste and emesis		1	
Bedpan	Containers for human waste and emesis	Fracture-type acceptable	1	Disposable or sanitized device
Urinal	Containers for human waste and emesis		1	Disposable or sanitized device
Blankets – Adult	covers with thermal insulating capabilities		2	Acrylic, wool, quilted or similar fabric
Baby Blanket with Head Covering	covers with thermal insulating capabilities		1	Flannel or fleece preferred Silver Swaddler® <u>not</u> acceptable as a substitute
Sheets	For cot and patient covers		4	Cotton, linen or disposable material

OBSTETRICAL EMERGENCIES PACK	Prepared package or kit: with Drape towel or under pad; Gauze dressings, Sterile gloves; Bulb syringe or aspirator; Cord clamps and/or umbilical ties; Plastic bags and ties for placental tissues; Infant receiving blanket or swaddling materials with a head covering.		1 Pack	
------------------------------	---	--	--------	--

INFECTION CONTROL SUPPLIES	Personal protective equipment to include: Disposable gloves sized for the crew; Fluid proof gowns or lab coats; Two face masks (NIOSH approved to at least N-95 standards); Eye shields or protective face shields; and Protective footwear or shoe covers.		Boxes of <u>non-latex gloves</u> sized for crew members; At least two each of other supplies or two commercial packages of isolation items	
Red plastic bags or trash bags labeled for biohazard	For infectious or biologically contaminated trash	at least 24" by 30"	2 bags	

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
Sharps disposal supplies	A puncture resistant container shall be supplied for sharps disposal in a locking-style bracket or in a locked compartment within the ambulance; a sheath style shall be supplied for on-scene use		1 installed 1 portable	See Rule text and TOSHA/CDC Requirements for Blood-borne Pathogens

#### INTRAVENOUS THERAPY SUPPLIES

Macro drip fluid adm. sets	ten to twenty drops per milliliter		3	
Micro drip fluid adm. sets	sixty drops per milliliter		3	
Antiseptic wipes	alcohol or approved antiseptic		12	
I.V. Catheters	over-the-needle type	14, 16,18,20,22 and 24 gauge	4 sets in each size (24)	
Intravenous solutions	Three liters of intravenous solutions, two liters of which will be crystalloid fluids	Containers May be in increments of 250, 500, or 1,000 mL	3,000 mL	Minimum of 2,000 mL of saline or LRS; Plastic containers required
Venous tourniquets	Disposable (non-latex) venous tourniquets, sufficient for adult and pediatric use.		2	
Intraosseous infusion needles		minimum 18 gauge	2	May be incorporated in Board approved devices

#### CARDIAC DEFIBRILLATOR/MONITORS

Cardiac monitor, electrocardiographic recorder, and defibrillator shall be provided for each ALS Ambulance		The ALS defibrillator shall provide a minimum setting of ten (10) joules.	1 Device Cardiac monitoring leads: electrodes Six for adults; Six for pediatric patients; Spare batteries	A biphasic waveform shall be required on any cardiac monitor/ defibrillator purchased for ambulances after the effective date of this rule
<b>AUTOMATED EXTERNAL DEFIBRILLATOR</b>	An AED shall be provided on each staffed ambulance, except those otherwise staffed and equipped to provide advanced life support		1 Device; 2 sets adult pads, and pediatric pads where capable with device. Spare batteries as per manufacturer	A biphasic waveform shall be required on any defibrillator purchased for ambulances after the effective date of this rule.

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
<b>MEDICATIONS AND DRUGS For Basic Ambulances</b>				
For treatment of Anaphylaxis	Epinephrine 1:1,000 preloaded syringe or a Tuberculin syringe with a minimum 5/8 inch, 25 gauge needle	0.3 mL per dose, sufficient quantity of Epinephrine 1:1,000 to administer two (2) doses to two patients	4 doses	
For treatment of Suspected Cardiac Patients	Aspirin, U.S.P.	81 mg or 325 mg	Sufficient for two doses	As indicated by service standing orders or protocol
For treatment of acute pulmonary distress	Beta-adrenergic agonist (albuterol, etc.)	Quantity sufficient for two repeated treatments	Nebulized treatments with appropriate administration devices	Or therapeutic equivalent identified by medical director for acute pulmonary distress.
For treatment of Chest Pain or Suspected Cardiac Event	Nitroglycerine	1/150 grain (0.4 mg)	bottle of thirty (30) tablets or sublingual spray	Or therapeutic equivalent identified by medical director

<b>MEDICATIONS AND DRUGS For Advanced Life Support Ambulances</b>	Medications used on advanced level ambulances shall be compatible with current standards as indicated by the American Heart Association's Emergency Cardiovascular Care Committee			
Cardiovascular medications	Adenosine	6 mg/2mL	Sufficient for doses up to 18 milligrams	Or therapeutic equivalent identified by medical director
	Atropine sulfate	1.0mg/10mL	four (4) prefilled syringes	
	Antiarrhythmic agents			
	Amiodarone	150 to 300 mg	to total at least 450 mg	
	Or Lidocaine	100 mg. in 5 mL	at least four prefilled syringes	Admixtures or premixed solutions shall be provided for maintenance drips
	Magnesium Sulfate	1 gram	Sufficient to administer 2 gm	
Bacteriostatic water and sodium chloride	For injection and dilution of medications			
Analgesics	Morphine, meperidine hydrochloride, nalbuphine (Nubain), butorphanol (Stadol), Nitrous oxide	As identified by EMS Medical Director and consigned to unit		Or therapeutic equivalent identified by medical director
Benzodiazepine anticonvulsant	Diazepam or other benzodiazepine in equivalent amounts sufficient to administer two successive maximum doses	Ten (10) milligrams/2mL	2 vials or prefilled syringes	Or therapeutic equivalent identified by medical director
Vasopressor agents	Epinephrine	1:10,000 of 1.0 mg/mL	4 prefilled syringes	Or therapeutic equivalent identified by medical director

ITEM	ITEM DESCRIPTION	MINIMUM SIZE/DOSE	NUMBER	COMMENTS Interpretive Guidance
Hypoglycemic countermeasures	Glucose testing devices for semi-quantitative blood glucose determinations		With device, media, calibration strips, and lancets	
Dextrose 50% in water		25 grams in 50 milliliters	2 prefilled syringes	Or therapeutic equivalent identified by medical director
Dextrose 25% in water		12.5 grams in 50 milliliters	2 prefilled syringes	Or therapeutic equivalent identified by medical director
Narcotic antagonist	Narcan (naloxone)	1mg/mL	2 ampules or prefilled syringes	Or therapeutic equivalent identified by medical director
Alkalinizing agents	Sodium Bicarbonate	50 mEq in 50 milliliters	2 ampules or prefilled syringes	Or therapeutic equivalent identified by medical director
Systemic diuretics	Furosemide	10 mg/mL	ampules, vials, or prefilled syringes to total 80 milligrams	Or therapeutic equivalent identified by medical director
Antinauseant	Promethazine IM Inject able	25mg/mL		Or therapeutic equivalent identified by medical director
Antihistamine	Diphenhydramine IM Inject able	50/mg		Or therapeutic equivalent identified by medical director
Syringes	for drug administration	1 mL, 3 mL, and 10 mL sizes	Assorted	With needles for administration
Pediatric Length - Based Resuscitation Tape	2002 Broselow™ or successor edition		1	